

Journal of Algebra and Its Applications
 Vol. 15, No. 8 (2015) 1650149 (18 pages)
 © World Scientific Publishing Company
 DOI: 10.1142/S0219498816501498



Intersections of quotient rings and Prüfer v -multiplication domains

Said El Baghdadi

*Department of Mathematics, Beni Mellal University
 23000 Beni Mellal, Morocco
 baghdadi@fstbm.ac.ma*

Marco Fontana

*Dipartimento di Matematica e Fisica
 Università degli Studi "Roma Tre", 00146 Rome, Italy
 fontana@mat.uniroma3.it*

Muhammad Zafrullah

*Department of Mathematics and Statistics
 Idaho State University, Pocatello, ID 83209-8085, USA
 mzafrullah@usa.net*

Received 11 August 2015

Accepted 14 September 2015

Published

Communicated by Roger Wiegand

Let D be an integral domain with quotient field K . Call an overring S of D a subring of K containing D as a subring. A family $\{S_\lambda \mid \lambda \in \Lambda\}$ of overrings of D is called a defining family of D , if $D = \bigcap \{S_\lambda \mid \lambda \in \Lambda\}$. Call an overring S a sublocalization of D , if S has a defining family consisting of rings of fractions of D . Sublocalizations and their intersections exhibit interesting examples of semistar or star operations [D. D. Anderson, Star operations induced by overrings, *Comm. Algebra* **16** (1988) 2535–2553]. We show as a consequence of our work that domains that are locally finite intersections of Prüfer v -multiplication (respectively, Mori) sublocalizations turn out to be Prüfer v -multiplication domains (PvMDs) (respectively, Mori); in particular, for the Mori domain case, we reobtain a special case of Théorème 1 of [J. Querré, Intersections d'anneaux entiers, *J. Algebra* **43** (1976) 55–60] and Proposition 3.2 of [N. Dessagnes, Intersections d'anneaux de Mori — exemples, *Port. Math.* **44** (1987) 379–392]. We also show that, more than the finite character of the defining family, it is the finite character of the star operation induced by the defining family that causes the interesting results. As a particular case of this theory, we provide a purely algebraic approach for characterizing PvMDs as a subclass of the class of essential domains (see also Theorem 2.4 of [C. A. Finocchiaro and

S. El Baghdadi, M. Fontana & M. Zafrullah

F. Tartarone, On a topological characterization of Prüfer v -multiplication domains among essential domains, preprint (2014), arXiv:1410.4037).

Keywords: Defining family of overrings; finite character; Mori, strong Mori and Krull domain; star and semistar operation; v -, τ -, w -operation; Prüfer and Dedekind domain; Prüfer v -multiplication domain.

Mathematics Subject Classification: 13F05, 13G05, 13B30, 13E99

1. Introduction

Throughout this paper D denotes an integral domain and K its quotient field.

A family of overrings (rings between D and K) $\{S_\lambda \mid \lambda \in \Lambda\}$ of D such that $D = \bigcap \{S_\lambda \mid \lambda \in \Lambda\}$ is called a *defining family of D* . We say that a *defining family is locally finite* (or, *has finite character*) if every nonzero element of D is a unit in all but a finite number of the S_λ 's.

When the rings S_λ are quotient rings of D , we get a representation of D as an intersection of quotient rings. This is the case of an important class of classical domains, e.g. the class of essential domains (definition recalled later), which includes Dedekind domains, Krull domains, Prüfer domains and their generalization Prüfer v -multiplication domains (PvMD, definition recalled later). A more general interesting representation is when each $S_\lambda \in \{S_\lambda \mid \lambda \in \Lambda\}$ is itself an intersection of quotient rings, e.g. if each S_λ is (τ) -flat over D (definition recalled later).

In this paper, we study some of these representations defined by appropriate finite character type conditions.

The theory of star and semistar operations is one of the key ingredients in achieving this goal. In fact, any representation associated to a defining family of a domain D induces a star operation on D (see [1]). More generally, any intersection of overrings of D defines a semistar operation on D (see for instance [11]). In this paper, we will mainly use the last more general setting.

The aim of this paper is to shed new light on some questions in the literature related to representations of domains as intersections of quotient rings. For instance, a well-known result in this area is a simple and elegant characterization, in the τ -finite character case, of the PvMDs given by Griffin in [23]: they are exactly the essential domains. An example by Heinzer and Ohm [25] shows that there exist essential domains that are not PvMDs. The question of when an essential domain is a PvMD was solved recently in [9], by using topological methods. In this paper, we introduce a weak form of the finite character property of a defining family of a domain which turns out to be the key idea for an algebra theoretic proof of the question of when an essential domain is a PvMD.

In Secs. 2 and 3, we give an overview on the theory of semistar operations and its interaction with a representation of a domain as an intersection of overrings. In Sec. 4, we investigate the question of when an intersection of a family of PvMDs is a PvMD. In the case of an intersection of overrings with finite character, we give an affirmative answer to the previous question, providing a generalization of a similar

well-known fact concerning the Krull domains, i.e. a locally finite intersection of Krull domains is a Krull domain. In Sec. 5, we provide a purely algebraic approach for characterizing PvMDs as a subclass of the class of essential domains.

2. Preliminaires

Throughout this paper, let D be an integral domain with quotient field K . Let $\overline{\mathbf{F}}(D)$ (respectively, $\mathbf{F}(D)$; $\mathbf{f}(D)$) be the set of all nonzero D -submodules of K (respectively, nonzero fractional ideals; nonzero finitely generated fractional ideals) of D (thus, $\mathbf{f}(D) \subseteq \mathbf{F}(D) \subseteq \overline{\mathbf{F}}(D)$).

A mapping $\star : \overline{\mathbf{F}}(D) \rightarrow \overline{\mathbf{F}}(D)$, $E \mapsto E^\star$, is called a *semistar operation* of D if, for all $z \in K$, $z \neq 0$ and for all $E, F \in \overline{\mathbf{F}}(D)$, the following properties hold: (\star_1) $(zE)^\star = zE^\star$; (\star_2) $E \subseteq F \Rightarrow E^\star \subseteq F^\star$; (\star_3) $E \subseteq E^\star$; and (\star_4) $E^{\star\star} := (E^\star)^\star = E^\star$.

When $D^\star = D$, \star is called a *(semi)star operation* on D ; in this case, the restriction of \star to $\mathbf{F}(D)$ is a usual *star operation* (see [20, Sec. 32] for more details).

As in the classical star-operation setting, we associate to a semistar operation \star of D a new semistar operation \star_f of D by setting, for every $E \in \overline{\mathbf{F}}(D)$,

$$E^{\star_f} := \bigcup \{F^\star \mid F \subseteq E, F \in \mathbf{f}(D)\}.$$

We call \star_f the semistar operation of finite type of D associated to \star . If $\star = \star_f$, we say that \star is a *semistar operation of finite type* on D . Note that $(\star_f)_f = \star_f$, so \star_f is a semistar operation of finite type of D .

We denote by $\mathbf{SStar}(D)$ (respectively, $\mathbf{SStar}_f(D)$) the set of all semistar operations (respectively, semistar operations of finite type) on D . Given two semistar operations \star' and \star'' of D , we say that $\star' \leq \star''$ if $E^{\star'} \subseteq E^{\star''}$, for all $E \in \overline{\mathbf{F}}(D)$. The relation “ \leq ” introduces a partial ordering in $\mathbf{SStar}(D)$. From the definition of \star_f , we deduce that $\star_f \leq \star$ and that \star_f is the largest semistar operation of finite type smaller than or equal to \star .

A semistar operation \star defined on an integral domain D is called *stable* provided that, for any $E, H \in \overline{\mathbf{F}}(D)$, we have $(E \cap H)^\star = E^\star \cap H^\star$. We denote by $\overline{\mathbf{SStar}}(D)$ the set of stable semistar operations on D .

Given a semistar operation \star on D , we can always associate to \star a stable semistar operation $\overline{\star}$ by defining, for every $E \in \overline{\mathbf{F}}(D)$,

$$E^{\overline{\star}} := \bigcup \{(E : I) \mid I \text{ nonzero ideal of } D \text{ such that } I^\star = D^\star\}.$$

It is easy to see that $\overline{\star} \leq \star$ and, moreover, that $\overline{\star}$ is the largest stable semistar operation that precedes \star . Therefore, \star is stable if and only if $\star = \overline{\star}$ [11, Proposition 3.7, Corollary 3.9].

As in the case of $\overline{\star}$, we can associate to each semistar operation \star a stable semistar operation of finite type $\widetilde{\star}$ by defining, for every $E \in \overline{\mathbf{F}}(D)$,

$$E^{\widetilde{\star}} := \bigcup \{(E : J) \mid J \text{ nonzero finitely generated ideal of } D \text{ such that } J^\star = D^\star\}.$$

S. El Baghdadi, M. Fontana & M. Zafrullah

The stable *semistar operation* of finite type $\tilde{\star}$ is smaller than or equal to \star , and it is the biggest stable *semistar operation* of finite type smaller than or equal to \star . It follows that \star is stable of finite type if and only if $\star = \tilde{\star}$. We denote by $\widetilde{\text{SStar}}(D)$ the set of stable *semistar operations* of finite type on D .

Let $\mathcal{S} := \{S_\lambda \mid \lambda \in \Lambda\}$ be a nonempty family of overrings of an integral domain D . Let $\wedge_{\mathcal{S}}$ be the *semistar operation* on D defined, for each $E \in \overline{\mathbf{F}}(D)$, by:

$$E^{\wedge_{\mathcal{S}}} := \bigcap \{ES_\lambda \mid \lambda \in \Lambda\}.$$

In particular, if S is an overring of D and $\mathcal{S} := \{S\}$, then the operation $\wedge_{\{S\}}$ is a semistar operation of finite type. If S is a D -flat overring, then $\wedge_{\{S\}}$ is a *stable semistar operation* (and of finite type) and conversely (see [32, Theorem 7.4(i); 37, Proposition 1.7]). In general, for each nonempty family \mathcal{S} of D -flat overrings of D , $\wedge_{\mathcal{S}}$ is stable, but it is not necessarily of finite type.

If Y is a nonempty subset of the prime spectrum $\text{Spec}(D)$ of an integral domain D , then we define the *semistar operation* \mathfrak{s}_Y induced by Y as the *semistar operation* associated to the set $\mathcal{T}(Y) := \{D_P \mid P \in Y\}$, i.e. $\mathfrak{s}_Y := \wedge_{\mathcal{T}(Y)}$ is the semistar operation defined by

$$E^{\mathfrak{s}_Y} := \bigcap \{ED_P \mid P \in Y\}, \text{ for every } E \in \overline{\mathbf{F}}(D).$$

A *semistar operation* of the type \mathfrak{s}_Y , for some $Y \subseteq \text{Spec}(D)$, is called a *spectral semistar operation* on D . We denote by $\text{SStar}_{\text{sp}}(D)$ the set of spectral *semistar operations* on D . Clearly, a spectral *semistar operation* is stable, i.e. $\text{SStar}_{\text{sp}}(D) \subseteq \widetilde{\text{SStar}}(D)$. Moreover, it is known that the previous sets of *semistar operations* coincide in the finite type case (see for instance [34, Lemma 1.32]):

$$\text{SStar}_{\text{sp}}(D) \cap \text{SStar}_f(D) = \overline{\text{SStar}}(D) \cap \text{SStar}_f(D) = \widetilde{\text{SStar}}(D).$$

For star operations \star , the notion of a “star-ideal” (that is, a nonzero ideal I of D , such that $I^\star = I$) is very useful. For a *semistar operation* \star , we need a more general notion, that coincides with the notion of star-ideal, when \star is a (semi)star operation. We say that a nonzero ideal I of D is a quasi- \star -ideal if $I^\star \cap D = I$. For example, it is easy to see that, for each nonzero ideal I of D such that $I^\star \cap D \neq D$, then $J := I^\star \cap D$ is a quasi- \star -ideal of D that contains I ; in particular, a \star -ideal (i.e. a nonzero ideal I such that $I^\star = I$) is a quasi- \star -ideal. Note that $I^\star \cap D \neq D$ is equivalent to $I^\star \neq D^\star$. A *quasi- \star -prime* is a quasi- \star -ideal which is also a prime ideal. We call a *quasi- \star -maximal* a maximal element in the set of all proper quasi- \star -ideals of D . We denote by $\text{QSpec}^\star(D)$ (respectively, $\text{QMax}^\star(D)$) the set of all quasi- \star -primes (respectively, quasi- \star -maximals) of D . It is well known that a quasi- \star -maximal ideal is a prime ideal and it is possible to prove that each quasi- \star_f -ideal is contained in a quasi- \star_f -maximal ideal (see for instance [14, Lemma 2.3]). When \star is a (semi)star operation, we simply set $\text{Max}^\star(D)$ (respectively, $\text{Spec}^\star(D)$) instead of $\text{QMax}^\star(D)$ (respectively, $\text{QSpec}^\star(D)$).

A *semistar operation* \star on an integral domain D is said to be an *eab semistar operation* (respectively, an *ab semistar operation*) if, for every $F, G, H \in \mathbf{f}(D)$

(respectively, for every $F \in \mathbf{f}(D)$, $G, H \in \overline{\mathbf{F}}(D)$) the inclusion $(FG)^* \subseteq (FH)^*$ implies $G^* \subseteq H^*$. Note that, if \star is **eab**, then \star_f is also **eab**, since \star and \star_f agree on nonzero finitely generated fractional ideals. We can associate to any semistar operation \star of D an **eab** semistar operation of finite type \star_a of D , called *the eab semistar operation associated to \star* , defined as follows for each $F \in \mathbf{f}(D)$ and for each $E \in \overline{\mathbf{F}}(D)$:

$$F^{\star_a} := \bigcup \{((FH)^* : H^*) \mid H \in \mathbf{f}(D)\},$$

$$E^{\star_a} := \bigcup \{F^{\star_a} \mid F \subseteq E, F \in \mathbf{f}(D)\},$$

[13, Definition 4.4, Proposition 4.5] (cf. also [15] and [16]). The previous construction, in the ideal systems setting, is essentially due to Jaffard [28] and Halter-Koch [24].

Obviously $(\star_f)_a = \star_a$. Note also that, when $\star = \star_f$, then \star is **eab** if and only if $\star = \star_a$ [13, Proposition 4.5(5)].

A *valuative semistar operation* is a semistar operation of the type $\wedge_{\mathcal{V}}$, where \mathcal{V} is a family of valuation overrings of D ; it is easy to see that $\wedge_{\mathcal{V}}$ is an **eab** semistar operation. In particular, if \mathcal{V} is the set of all valuation overrings of D , the **b**-operation, where $\mathbf{b} := \wedge_{\mathcal{V}}$, is an **eab** semistar operation of finite type on D (see [8, Proposition 4.5; 20, pp. 394 and 398]).

Just as in the case of the relation between stable and spectral operations, not every **eab** semistar operation is valutive, but the two definitions agree on finite type operations (see, for instance, [13, Corollaries 3.8 and 5.2]).

Denote by $\mathbf{SStar}_{\text{val}}(D)$ (respectively, $\mathbf{SStar}_{\text{eab}}(D)$; $\mathbf{SStar}_{f, \text{eab}}(D)$) the set of valutive (respectively, **eab**; **eab** of finite type) semistar operations on D . By the previous remarks, we have:

$$\mathbf{SStar}_{f, \text{eab}}(D) := \mathbf{SStar}_{\text{eab}}(D) \cap \mathbf{SStar}_f(D) = \mathbf{SStar}_{\text{val}}(D) \cap \mathbf{SStar}_f(D).$$

3. Sublocalizations and Associated Semistar Operations

Let D be an integral domain and S an overring of D . It is possible to define an “extension” map $\text{ext} := \text{ext}(D, S) : \mathbf{SStar}(D) \rightarrow \mathbf{SStar}(S)$ (respectively, “contraction” map $\text{con} := \text{con}(S, D) : \mathbf{SStar}(S) \rightarrow \mathbf{SStar}(D)$), by setting $\star \mapsto \star^e$ (respectively, $\star \mapsto \star^c$), where:

$$\star^e : \overline{\mathbf{F}}(S) \subseteq \overline{\mathbf{F}}(D) \xrightarrow{\star} \overline{\mathbf{F}}(D) \xrightarrow{\otimes_D S} \overline{\mathbf{F}}(S), \quad F \mapsto (F^*)S$$

$$\text{(respectively, } \star^c : \overline{\mathbf{F}}(D) \xrightarrow{\otimes_D S} \overline{\mathbf{F}}(S) \xrightarrow{\star} \overline{\mathbf{F}}(S) \subseteq \overline{\mathbf{F}}(D), \quad E \mapsto (ES)^*).$$

Note that $(F^*)S = F^* \in \overline{\mathbf{F}}(S)$, since for each nonzero $s \in S$, $sF^* = (sF)^* \subseteq F^*$, being $F \in \overline{\mathbf{F}}(S)$.

We collect in the following lemma some basic properties of the maps $\text{ext}(D, S)$ and $\text{con}(S, D)$ (see also, for instance, [34, Proposition 1.35, Lemma 1.36, Example 1.37, Propositions 2.11(1), 2.13(1) and 2.15]).

S. El Baghdadi, M. Fontana & M. Zafrullah

- Lemma 3.1.** (1) *The map \mathbf{ext} is order-preserving, i.e. $\star_1 \leq \star_2$ implies $(\star_1)^\circ \leq (\star_2)^\circ$.*
(2) *The map \mathbf{ext} preserves semistar operations of finite type, i.e. $\mathbf{ext}|_{\mathbf{SStar}_f(D)} : \mathbf{SStar}_f(D) \rightarrow \mathbf{SStar}_f(S)$.*
(3) *The map \mathbf{con} is order-preserving, i.e. $\star_1 \leq \star_2$ implies $(\star_1)^\circ \leq (\star_2)^\circ$.*
(4) *The map \mathbf{con} preserves semistar operations of finite type, i.e. $\mathbf{con}|_{\mathbf{SStar}_f(S)} : \mathbf{SStar}_f(S) \rightarrow \mathbf{SStar}_f(D)$.*
(5) *Let \mathbf{d}_D (respectively, \mathbf{d}_S) the identity semistar operation on D (respectively, on S), then $(\mathbf{d}_D)^\circ = (\wedge_{\{S\}})^\circ = \mathbf{d}_S$.*
(6) $(\mathbf{d}_S)^\circ = \wedge_{\{S\}}$.
(7) *For each $\star \in \mathbf{SStar}(S)$, $(\star^\circ)^\circ = \star$ (i.e. $\mathbf{ext} \circ \mathbf{con} = \mathbf{id}_{\mathbf{SStar}(S)}$).*
(8) *For each $\star \in \mathbf{SStar}(D)$, $(\star^\circ)^\circ \geq \star$ (for short, we summarize this property by writing $\mathbf{con} \circ \mathbf{ext} \geq \mathbf{id}_{\mathbf{SStar}(D)}$). In particular, if $D \subsetneq S$, $\mathbf{d}_D \leq ((\mathbf{d}_D)^\circ)^\circ = (\mathbf{d}_S)^\circ = \wedge_{\{S\}}$.*

Remark 3.2. In relation with statements (2) and (4) of the previous lemma, we observe that \mathbf{ext} preserves stable semistar operations and, if $D^\star = S$, then \star is spectral on D implies that $(\star)^\circ$ is spectral on S [34, Proposition 2.11(2) and (6)]. On the other hand, \mathbf{con} preserves neither stability nor spectrality. For instance, \mathbf{d}_S is obviously spectral and hence stable on S while, if S is not a D -flat overring of D , $(\mathbf{d}_S)^\circ = \wedge_{\{S\}}$ is not stable (and, a fortiori, is not spectral) on D .

The overring S of D is a *sublocalization* of D if S is a nonempty intersection of ring of fractions of D . Thus S is a sublocalization of D if and only if there exists a nonempty family $\{T_\alpha \mid \alpha \in \mathcal{A}\}$ of multiplicatively closed subsets of nonzero elements of D such that $S = \bigcap \{D_{T_\alpha} \mid \alpha \in \mathcal{A}\}$. It is well known that a sublocalization S of D is an intersection of localizations of D at prime ideals, since each ring of fractions of D is an intersection of localizations of D (see [21, 36]). Indeed if T is a multiplicatively closed subset of an integral domain D , with $0 \notin T$, then $D_T = \bigcap \{D_P \mid P \in \text{Spec}(D) \text{ and } P \cap T = \emptyset\}$. Therefore, if $\{T_\alpha \mid \alpha \in \mathcal{A}\}$ is a family of multiplicatively closed sets of nonzero elements of D and $S = \bigcap \{D_{T_\alpha} \mid \alpha \in \mathcal{A}\}$, then $S = \bigcap \{D_P \mid P \in \text{Spec}(D), P \cap T_\alpha = \emptyset, \text{ for some } \alpha \in \mathcal{A}\}$.

From the previous remarks, we deduce immediately the following.

Lemma 3.3. *Let S be an overring of D . Then, S is a sublocalization of D if and only if $S = \bigcap \{D_P \mid P \in \text{Spec}(D), S \subseteq D_P\}$.*

Recall that, by [36, Theorem 1], S is a D -flat overring of D if and only if, for each $P \in \text{Spec}(D)$, either $PS = S$ or $S \subseteq D_P$. Therefore, by Lemma 3.3, if S is a D -flat overring of D then S is a sublocalization of D . However, the converse is not true [26, Sec. 2, Discussion 2.1].

Proposition 3.4. *Let D be a domain and let $S = \bigcap \{D_{T_\alpha} \mid \alpha \in \mathcal{A}\}$ be a sublocalization of D , where $\{T_\alpha \mid \alpha \in \mathcal{A}\}$ is a given family of multiplicatively closed subsets of nonzero elements of D . Set $\mathcal{T} := \mathcal{T}(S) := \{D_{T_\alpha} \mid \alpha \in \mathcal{A}\}$, considered as a family of*

overrings of S , let $* := \wedge_{\mathcal{T}(S)} \in \text{SStar}(S)$ and set $\star_S := *^c = * \circ \wedge_{\{S\}} \in \text{SStar}(D)$, i.e. $E^{\star_S} := (ES)^* = \cap\{ED_{T_\alpha} \mid \alpha \in \mathcal{A}\}$, for each $E \in \overline{\mathbf{F}}(D)$.

- (1) $\star_S = \wedge_{\mathcal{T}(D)}$, where in the last equality the family $\mathcal{T}(D)$ is the family \mathcal{T} considered as a family of overrings of D .
- (2) $\widetilde{\wedge_{\{S\}}} \leq \wedge_{\{S\}} \leq \star_S$.
- (3) Let $A_1, A_2, \dots, A_n \in \overline{\mathbf{F}}(D)$. Then, $(A_1 \cap A_2 \cap \dots \cap A_n)^{\star_S} = ((A_1 \cap A_2 \cap \dots \cap A_n)S)^* = (A_1S)^* \cap (A_2S)^* \cap \dots \cap (A_nS)^* = A_1^{\star_S} \cap A_2^{\star_S} \cap \dots \cap A_n^{\star_S}$, i.e. \star_S is a stable semistar operation on D .
- (4) If $F \in \mathbf{f}(D)$, then $(FS)^{-1} = (F^{-1}S)^* = (F^{-1})^{\star_S}$.
- (5) Let $\mathbf{v}(S)$ be the (semi)star \mathbf{v} -operation of S (i.e. $E^{\mathbf{v}(S)} := (S : (S : E))$) for each $E \in \overline{\mathbf{F}}(D)$). If $F \in \mathbf{f}(D)$, then $(FS)^{-1} = (F^{-1}S)^{\mathbf{v}(S)}$.

Proof. (1) is a straightforward consequence of the definitions.

(2) In general, for each semistar operation \star on D , the stable semistar operation of finite type $\tilde{\star}$ is such that $\tilde{\star} \leq \star$. The second inequality follows by observing that $ES \subseteq (ES)^*$, for each $E \in \overline{\mathbf{F}}(D)$.

(3) follows easily from the fact that \star_S coincides with $(\wedge_{\mathcal{T}(S)})^c$ and $\mathcal{T}(S)$ is a family of overrings of fractions of S (and D), hence $* = \wedge_{\mathcal{T}(S)}$ (respectively, $(\wedge_{\mathcal{T}(S)})^c$) is a stable semistar operation on S (respectively, on D).

(4) Let $F = (f_1, f_2, \dots, f_r)$, then $(FS)^{-1} = (S : FS) = \cap\{f_i^{-1}S \mid 1 \leq i \leq r\}$ and $F^{-1} = (D : F) = \cap\{f_i^{-1}D \mid 1 \leq i \leq r\}$. Therefore, $(F^{-1})^{\star_S} = (\cap\{f_i^{-1}D \mid 1 \leq i \leq r\})^{\star_S} = \cap\{(f_i^{-1}D)^{\star_S} \mid 1 \leq i \leq r\} = \cap\{(f_i^{-1}S)^* \mid 1 \leq i \leq r\} = \{f_i^{-1}S \mid 1 \leq i \leq r\} = (FS)^{-1}$. Thus $(F^{-1}S)^* = (F^{-1})^{\star_S} = (FS)^{-1}$.

(5) Since $*$ is a (semi)star operation of S , it is clear that $* \leq \mathbf{v}(S)$ (see [20, Theorem 34.1(4); 34, Lemma 1.11]). By (4), we have $(FS)^{-1} = (F^{-1}S)^*$. Therefore, $((FS)^{-1})^{\mathbf{v}(S)} = (FS)^{-1} = (F^{-1}S)^* = ((F^{-1}S)^*)^{\mathbf{v}(S)} = (F^{-1}S)^{\mathbf{v}(S)}$. \square

Remark 3.5. As a straightforward consequence of the previous proposition, we re-obtain the following well-known properties. If S is a D -flat overring of D , then

- (1) for $A_1, A_2, \dots, A_n \in \overline{\mathbf{F}}(D)$, $(A_1 \cap A_2 \cap \dots \cap A_n)S = A_1S \cap A_2S \cap \dots \cap A_nS$;
- (2) for each $F \in \mathbf{f}(D)$, $(FS)^{-1} = F^{-1}S$ and $(FS)^{\mathbf{v}(S)} = (F^{\mathbf{v}(D)}S)^{\mathbf{v}(S)}$, where $\mathbf{v}(S)$ (respectively, $\mathbf{v}(D)$) is the (semi)star \mathbf{v} -operation of S (respectively, of D), for details see [10, Proposition 0.6(b)].

Let \star be a semistar operation on the integral domain D . For $E \in \overline{\mathbf{F}}(D)$, we say that E is \star -finite if there exists a $F \in \mathbf{f}(D)$ such that $F^\star = E^\star$. (Note that in the above definition, we do not require that $F \subseteq E$.) It is immediate to see that if $\star_1 \leq \star_2$ are semistar operations and E is \star_1 -finite, then E is \star_2 -finite. In particular, if E is \star_j -finite, then it is \star -finite. The converse is not true in general (see [17, Remark 2.4]), and one can prove that E is \star_j -finite if and only if there exists $F \in \mathbf{f}(D)$, $F \subseteq E$, such that $F^\star = E^\star$ [17, Lemma 2.3]. This result was proved in the star operation setting by Zafrullah in [40, Theorem 1.1].

S. El Baghdadi, M. Fontana & M. Zafrullah

Lemma 3.6. *Let S be an overring of D and $*$ a semistar operation on S . Consider the semistar operation $\star_S := {}^c$ on D . Let I be a nonzero ideal of D and assume that $I^{\star_S} := (IS)^* = ((x_1, x_2, \dots, x_n)S)^*$, where $x_k \in IS$, for $1 \leq k \leq n$. Then, we can find a finitely generated ideal J of D , with $J \subseteq I$, such that,*

$$I^{\star_S} = (IS)^* = (JS)^* = J^{\star_S}.$$

Proof. Indeed, as $x_k \in IS$, we have $x_k = \sum_{j=1}^{r_k} i_{kj}s_j$, where $i_{kj} \in I$ and $s_j \in S$. Then $x_k S \subseteq I_k S$ for some finitely generated ideal $I_k \subseteq I$ of D , for every k . Take $J := \sum_k I_k$ and the verification of the claim is straightforward. \square

Proposition 3.7. *Let $\{S_\lambda \mid \lambda \in \Lambda\}$ be a family of overrings of D and let $*_\lambda$ be a (semi)star operation on S_λ . Set $\star_{S_\lambda} := (*_\lambda)^c$, i.e. $E^{\star_{S_\lambda}} := (ES_\lambda)^{*\lambda}$, for each $E \in \overline{\mathbf{F}}(D)$. Consider the semistar operation on D , $\star := \wedge \star_{S_\lambda} : \overline{\mathbf{F}}(D) \rightarrow \overline{\mathbf{F}}(D)$, defined by $E \mapsto \bigcap \{(ES_\lambda)^{*\lambda} \mid \lambda \in \Lambda\}$.*

Suppose that $D = \bigcap \{S_\lambda \mid \lambda \in \Lambda\}$ is locally finite. If I is a nonzero ideal of D such that $(IS_\lambda)^{\lambda} = ((x_{\lambda 1}, x_{\lambda 2}, \dots, x_{\lambda n_\lambda})S_\lambda)^{*\lambda}$ with $x_{\lambda \mu} \in IS_\lambda$, for each $\lambda \in \Lambda$ and $1 \leq \mu \leq n_\lambda$, then there is a finitely generated ideal $J \subseteq I$ in D such that $J^\star = I^\star$.*

Proof. Since $D = \bigcap \{S_\lambda \mid \lambda \in \Lambda\}$ is locally finite, we have $IS_k \neq S_k$ for at most a finite subset $\{S_k \mid 1 \leq k \leq n\}$. Now, take a nonzero element $j \in I$, for the same reason, j is a nonunit in only finitely many overrings S_λ and, by the previous considerations, we can assume that j is a nonunit in $\{S_1, S_2, \dots, S_n, S_{n+1}, \dots, S_m \mid \text{for some } m \geq n\}$.

If $m = n$, then $jD \subseteq I$ is such that $jS_k \neq S_k$, precisely for $1 \leq k \leq n$.

If $m > n$, since $IS_h = S_h$ for each $n+1 \leq h \leq m$, there exists a finitely generated ideal $I_h \subseteq I$ such that $I_h S_h = S_h$. Thus, the finitely generated ideal $J_0 := jD + \sum_h I_h \subseteq I$ ensures that $J_0 S_k \neq S_k$, precisely for $1 \leq k \leq n$.

From Lemma 3.6, for each $\lambda \in \Lambda$, we know that $I^{\star_{S_\lambda}} = (IS_\lambda)^{*\lambda} = (J_\lambda S_\lambda)^{*\lambda} = (J_\lambda)^{\star_{S_\lambda}}$, for some finitely generated ideal $J_\lambda \subseteq I$ of D . In particular, if we consider the finite subset $\{S_k \mid 1 \leq k \leq n\}$ of $\{S_\lambda \mid \lambda \in \Lambda\}$, then we can find a finite set of finitely generated ideals $\{J_k \mid 1 \leq k \leq n\}$ contained in I such that $(IS_k)^{*\lambda} = (J_k S_k)^{*\lambda}$, where $*_k := *_{\lambda_k}$, for $1 \leq k \leq n$. Set $J := J_0 + J_1 + \dots + J_n$, by construction, it is easy to see that J is finitely generated ideal of D contained in I .

Therefore, $(IS_\lambda)^{*\lambda} = S_\lambda = (JS_\lambda)^{*\lambda}$, for each $\lambda \in \Lambda \setminus \{1, 2, \dots, n\}$ (since, in the present situation, $J_0 S_\lambda = JS_\lambda = IS_\lambda = S_\lambda$). For $k \in \{1, 2, \dots, n\}$, we have $(IS_k)^{*\lambda} = (J_k S_k)^{*\lambda} \subseteq (JS_k)^{*\lambda}$. Thus, for all $\lambda \in \Lambda$, we have $(IS_\lambda)^{*\lambda} \subseteq (JS_\lambda)^{*\lambda}$ and so we conclude that $I^\star \subseteq J^\star$. The opposite inclusion is trivial, since $J \subseteq I$. \square

4. Sublocalizations and Prüfer v -multiplication Domains

Let \star be a semistar operation on an integral domain D . For a nonzero ideal I of D , we say that I is \star -invertible if $(II^{-1})^\star = D^\star$ (cf. also [31]). From the fact that $\text{QMax}^{\tilde{\star}}(D) = \text{QMax}^{\star_f}(D)$, it easily follows that an ideal I is $\tilde{\star}$ -invertible if and

only if I is \star_f -invertible (note that if \star is a semistar operation of finite type, then $(II^{-1})^\star = D^\star$ if and only if $II^{-1} \not\subseteq M$ for all $M \in \text{QMax}^\star(D)$). It is well known that if I is \star_f -invertible, then I and I^{-1} are both \star_f -finite [17, Proposition 2.6].

An integral domain D is called a *Prüfer \star -multiplication domain* (for short, $P\star MD$) if every nonzero finitely generated ideal of D is \star_f -invertible (cf. for instance [12]). Note that for $\star = *$ a star operation of finite type on D , $P\star MD$ s were introduced by Houston, Malik, and Mott in [27] as $*$ -multiplication domains. When $\star = \nu$, we have the classical notion of PvMD (cf. for instance [23, 29, 33]); when $\star = \mathbf{d}$, where \mathbf{d} denotes the identity (semi)star operation, we have the notion of Prüfer domain [20, Theorem 22.1]. For star operations $*$, the only $P\star MD$ s are the PvMDs and the Prüfer domains since in a $P\star MD$, $\star_f = \mathbf{t}$ (see [12, Proposition 3.4; 29, Theorem 3.5]).

Note that from the definition and from the previous observations, it immediately follows that the notions of $P\star MD$, $P\star_f MD$, and $P\tilde{\star} MD$ coincide.

As in the star case [3, Corollary 2.10], it is well known that, for each semistar operation \star , we have $\tilde{\star} = \bigwedge_{Q \in \text{QMax}^{\star_f}(D)}$, i.e. for each $E \in \overline{\mathbf{F}}(D)$,

$$E^{\tilde{\star}} = \bigcap \{ED_Q \mid Q \in \text{QMax}^{\star_f}(D)\}.$$

From this fact, it can be deduced that D is a $P\star MD$ if and only if D_Q is a valuation domain for each $Q \in \text{QMax}^{\star_f}(D)$ [12, Theorem 3.1].

Recall that an *essential valuation overring* V of an integral domain D is a valuation overring of D such that $V = D_P$ for some $P \in \text{Spec}(D)$; in this situation, P is called *essential prime*. A family of overrings $\{S_\lambda \mid \lambda \in \Lambda\}$ of D is said to be an *essential representation* (or, an *essential defining family*) of D , if $D = \bigcap \{S_\lambda \mid \lambda \in \Lambda\}$ and each S_λ is an essential valuation overring of D . An *essential domain* is an integral domain having an essential representation. A PvMD is always essential because D_Q is a valuation domain for each $Q \in \text{QMax}^\nu(D)$ [29, Theorem 3.2].

Theorem 4.1. *Let $\{S_\lambda \mid \lambda \in \Lambda\}$ be a family of sublocalizations of an integral domain D . Suppose that $D = \bigcap \{S_\lambda \mid \lambda \in \Lambda\}$ where the intersection is locally finite.*

(1) *Let $\mathcal{T}_\lambda := \mathcal{T}(S_\lambda) := \{D_{T_{\alpha_\lambda}} \mid \alpha_\lambda \in \mathcal{A}_\lambda\}$ be a defining family of S_λ and let \ast_λ be the (semi)star operation on S_λ induced by \mathcal{T}_λ , i.e. $\ast_\lambda := \bigwedge_{\mathcal{T}_\lambda}$. As in Proposition 3.7, set $\star := \bigwedge \{(\ast_\lambda)^c \mid \lambda \in \Lambda\}$.*

Assume that, for each $\lambda \in \Lambda$,

- (a) *\ast_λ is a (semi)star operation of finite type of S_λ , and*
- (b) *S_λ is a $P\ast_\lambda MD$,*

then D is a $P\star MD$, and so D is a $P\nu MD$.

(2) *Assume that each of S_λ is a $P\nu MD$, then D is a $P\nu MD$.*

Proof. (1) Recall that, given a semistar operation \star on an integral domain D , D is a $P\star MD$ if and only if $\tilde{\star}$ is a **eab** semistar operation, i.e. $\tilde{\star} = \star_a$ [12, Theorem 3.1].

S. El Baghdadi, M. Fontana & M. Zafrullah

We start by observing that, in the present situation, $*_\lambda$ is a stable (semi)star operation, because the family \mathcal{T}_λ consists of rings of fractions. Since we are assuming that $*_\lambda$ is of finite type of S_λ and S_λ is a $P*_\lambda$ MD, we have $*_\lambda = \widetilde{*}_\lambda = (*_\lambda)_a$ [12, Theorem 3.1].

Moreover, $\star = \bigwedge\{(*_\lambda)^{c_\lambda} \mid \lambda \in \Lambda\}$ is a (semi)star operation of finite type of D , since the intersection $D = \bigcap\{S_\lambda \mid \lambda \in \Lambda\}$ is locally finite and each $*_\lambda$ is of finite type (see [1, Theorem 2(4); 8, Corollary 2.9]). Clearly, \star is a stable and valuative (semi)star operation on D , because, in the present setting, each $(*_\lambda)^{c_\lambda} (= \text{con}(D, S_\lambda)(*_\lambda))$ is a stable (since it is induced by a family of rings of fractions of D) and valuative (semi)star operation on D (since $*_\lambda$ is valuative and the valuation overrings of S_λ are valuation overrings of D). We conclude that $\widetilde{\star} = \star$ is an **eab** (semi)star operation and so D is a $P\star$ MD. Since $\star \leq \mathfrak{v}$, then D is also a $P\mathfrak{v}$ MD.

(2) For each $\lambda \in \Lambda$, we take as defining family of S_λ the family of valuation overrings $\mathcal{T}_\lambda := \mathcal{T}(S_\lambda) := \{(S_\lambda)_{\mathfrak{q}_\lambda} \mid \mathfrak{q}_\lambda \in \mathcal{A}_\lambda := \text{Max}^\tau(S_\lambda)\}$. In the present situation, the (semi)star operation on S_λ associated to \mathcal{T}_λ , i.e. $*_\lambda = \bigwedge_{\mathcal{T}_\lambda}$, coincides with \mathfrak{w}_λ , that is the \mathfrak{w} -operation on S_λ . It is easy to see that the assumptions of (1) are satisfied (after recalling that a $P\mathfrak{v}$ MD coincides with a $P\mathfrak{w}$ MD) and so, if we denote by \star the (semi)star operation $\bigwedge\{(\mathfrak{w}_\lambda)^{c_\lambda} \mid \lambda \in \Lambda\}$, we can conclude by (1) that D is a $P\star$ MD. In particular, since $\star \leq \mathfrak{v}$, D is a $P\mathfrak{v}$ MD. \square

Recall that an overring S of an integral domain D is a *t-flat overring of D* if, for each maximal \mathfrak{t} -ideal M of S , $S_M = D_{(M \cap D)}$ [30].

Remark 4.2. Note that it is possible to give a direct and independent proof of Theorem 4.1(2) under the assumptions that $D = \bigcap\{S_\lambda \mid \lambda \in \Lambda\}$, the intersection is locally finite and each S_λ is \mathfrak{t} -flat.

By assumption,

$$D = \bigcap\{S_\lambda \mid \lambda \in \Lambda\} = \bigcap\left\{\bigcap\{(S_\lambda)_{\mathfrak{q}_\lambda} \mid \mathfrak{q}_\lambda \in \text{Max}^\tau(S_\lambda)\} \mid \lambda \in \Lambda\right\}$$

and the valuation overring $(S_\lambda)_{\mathfrak{q}_\lambda}$ is essential for D , for each $\lambda \in \Lambda$ and for each $\mathfrak{q}_\lambda \in \text{Max}^\tau(S_\lambda)$. Now, by [39, Lemma 8], an essential domain D is a $P\mathfrak{v}$ MD if and only if, for every pair of elements $a, b \in D \setminus \{0\}$, the ideal $aD \cap bD$ is a \mathfrak{v} -ideal of finite type.

As each S_λ is a $P\mathfrak{v}$ MD, $aS_\lambda \cap bS_\lambda$ is a \mathfrak{v}_λ -ideal of finite type, thus we can find $x_{\lambda 1}, x_{\lambda 2}, \dots, x_{\lambda n_\lambda} \in S_\lambda$ such that

$$aS_\lambda \cap bS_\lambda = (x_{\lambda 1}, x_{\lambda 2}, \dots, x_{\lambda n_\lambda})^{\mathfrak{v}_\lambda} = (x_{\lambda 1}, x_{\lambda 2}, \dots, x_{\lambda n_\lambda})^{\mathfrak{v}_\lambda} = (x_{\lambda 1}, x_{\lambda 2}, \dots, x_{\lambda n_\lambda})^{\mathfrak{v}_\lambda}$$

where \mathfrak{v}_λ , \mathfrak{t}_λ , and \mathfrak{w}_λ are the \mathfrak{v} -, the \mathfrak{t} -, and \mathfrak{w} -operation on the S_λ 's, and we already observed that the \mathfrak{t} -, and \mathfrak{w} -operation coincide on the $P\mathfrak{v}$ MD S_λ [29, Theorem 3.5] (via [41, Theorem 4.7]).

On the other hand, by Proposition 3.4(3), $((aD \cap bD)S_\lambda)^{\mathfrak{v}_\lambda} = (aS_\lambda)^{\mathfrak{v}_\lambda} \cap (bS_\lambda)^{\mathfrak{v}_\lambda} = aS_\lambda \cap bS_\lambda$, for each λ . Therefore,

$$(aS_\lambda \cap bS_\lambda)^{\mathfrak{v}_\lambda} = (x_{\lambda 1}, x_{\lambda 2}, \dots, x_{\lambda n_\lambda})^{\mathfrak{v}_\lambda}$$

and, necessarily, $x_{\lambda k} \in aS_{\lambda} \cap bS_{\lambda}$, for $1 \leq k \leq n_{\lambda}$. Let $I := aD \cap bD$, by Lemma 3.6, we can find a finitely generated ideal $J_{\lambda} := (j_{\lambda 1}, j_{\lambda 2}, \dots, j_{\lambda r_{\lambda}})D$, with $J_{\lambda} \subseteq I$, such that,

$$(IS_{\lambda})^{*\lambda} = ((j_{\lambda 1}, j_{\lambda 2}, \dots, j_{\lambda r_{\lambda}})S_{\lambda})^{*\lambda}.$$

Next, as $D = \bigcap \{S_{\lambda} \mid \lambda \in \Lambda\}$ is of finite character then, in particular, \star is a (semi)star operation on D . Moreover, by Proposition 3.7, there exists a finitely generated ideal J of D , with $J \subseteq I$, such that $J^{\star} = I^{\star}$. Since $\star \leq \mathfrak{v}$, then $J^{\mathfrak{v}} = (J^{\star})^{\mathfrak{v}} = (I^{\star})^{\mathfrak{v}} = I^{\mathfrak{v}} = aD \cap bD$.

From the previous Theorem 4.1, we deduce immediately the following two corollaries.

Corollary 4.3. *Let $\{S_{\lambda} \mid \lambda \in \Lambda\}$ be a family of essential valuation overrings of D such that $D = \bigcap \{S_{\lambda} \mid \lambda \in \Lambda\}$, where the intersection is locally finite. Then D is a PvMD.*

Corollary 4.4. *Let $\{S_{\lambda} \mid \lambda \in \Lambda\}$ be a family of sublocalizations of an integral domain D . Assume that $D = \bigcap \{S_{\lambda} \mid \lambda \in \Lambda\}$, where the intersection is locally finite. If each of S_{λ} is a Prüfer domain, then D is a PvMD.*

Let \star be a semistar operation on an integral domain D . We say that D is a \star -Noetherian domain if D has the ascending chain condition on quasi- \star -ideals. Note that the \mathfrak{d} -Noetherian domains are just the usual Noetherian domains and the notions of \mathfrak{v} -Noetherian (respectively, \mathfrak{w} -Noetherian) domain and *Mori* (respectively, *strong Mori*) domain coincide (cf. also [38]). Recall that, in the star case, the concept of *star Noetherian domain* has been introduced by Zafrullah [40] (see, also, for instance, [2, 19, 34]).

The following properties follow easily from the definitions (for more details, see for instance [34, Lemmas 4.16 and 4.18 and Corollary 4.19] or [7, Lemmas 3.1 and 3.3]).

- (1) If $\star_1 \leq \star_2$ are two semistar operations on D , then D is \star_1 -Noetherian implies that D is \star_2 -Noetherian; in particular, a Noetherian domain is a \star -Noetherian domain, for any semistar operation \star on D .
- (2) If \star is a (semi)star operation and if D is a \star -Noetherian domain, then D is a Mori domain.
- (3) D is \star -Noetherian if and only if, for each nonzero ideal I of D , there exists a nonzero finitely generated ideal J of D such that $J \subseteq I$ and $J^{\star} = I^{\star}$.
- (4) D is \star -Noetherian if and only if D is \star_f -Noetherian; in particular, the notions of \mathfrak{v} -Noetherian domain and \mathfrak{t} -Noetherian domain coincide with the notion of Mori domain.

The following proposition extends to the semistar setting a result obtained by Querré in 1976 (see Remark 4.7).

S. El Baghdadi, M. Fontana & M. Zafrullah

Proposition 4.5. *Let $\{S_\lambda \mid \lambda \in \Lambda\}$ be a family of overrings of D such that $D = \bigcap \{S_\lambda \mid \lambda \in \Lambda\}$ and the intersection is locally finite. Let $*_\lambda$ be a (semi)star operation on S_λ and consider the semistar operation on D , $\star := \bigwedge \{(*_\lambda)^{c_\lambda} \mid \lambda \in \Lambda\}$.*

- (1) *Assume that, for each $\lambda \in \Lambda$, S_λ is $*_\lambda$ -Noetherian, then D is \star -Noetherian.*
- (2) *Assume that, for each $\lambda \in \Lambda$, S_λ is $\widetilde{*}_\lambda$ -Noetherian and that the semistar operation $\bullet := \bigwedge \{(\widetilde{*}_\lambda)^{c_\lambda} \mid \lambda \in \Lambda\}$ on D is stable (e.g. when the S_λ 's are quotient rings of D), then D is $\widetilde{\star}$ -Noetherian.*

Proof. (1) Given a nonzero ideal I of D , since S_λ is $*_\lambda$ -Noetherian there exists a nonzero finitely generated ideal J_λ in S_λ such that $J_\lambda \subseteq IS_\lambda$ and $(IS_\lambda)^{*_\lambda} = (J_\lambda)^{*_\lambda}$. Since $D = \bigcap \{S_\lambda \mid \lambda \in \Lambda\}$ and the intersection is locally finite, by Lemma 3.6 and Proposition 3.7, we can assume that $(J_\lambda)^{*_\lambda} = (JS_\lambda)^{*_\lambda}$, for each $\lambda \in \Lambda$, where J is a finitely generated ideal of D such that $J \subseteq I$ and $J^\star = I^\star$.

(2) Note that $\bullet \leq \widetilde{\star} \leq \star_f \leq \star$. Indeed, we have \bullet is stable; moreover \bullet is a (semi)star operation of finite type, since $(\widetilde{*}_\lambda)^{c_\lambda}$ is of finite type, for each $\lambda \in \Lambda$, and $D = \bigcap \{S_\lambda \mid \lambda \in \Lambda\}$ is locally finite (see [1, [Theorem 2(4); 8, [Proposition 2.9]]]).

If we show that D is \bullet -Noetherian, then *a fortiori* we have that D is $\widetilde{\star}$ -Noetherian. For this, given ideal I of D we have $(IS_\lambda)^{\widetilde{\star}} = \bigcap \{(IS_\lambda)_{\mathfrak{q}_{\lambda\mu}} \mid \mathfrak{q}_{\lambda\mu} \in \text{Max}^{\widetilde{\star}}(S_\lambda)\}$. Since S_λ is $\widetilde{*}_\lambda$ -Noetherian (and $\widetilde{*}_\lambda$ is of finite type), there exists a finitely generated ideal J_λ in S_λ such that $J_\lambda \subseteq IS_\lambda$ and $(J_\lambda)^{\widetilde{\star}} = (IS_\lambda)^{\widetilde{\star}}$. Again, as $D = \bigcap \{S_\lambda \mid \lambda \in \Lambda\}$ is locally finite, Proposition 3.7 applies, and so there exists a finitely generated ideal J of D , such that $J \subseteq I$ and $J^\bullet = I^\bullet$. Therefore, D is \bullet -Noetherian. \square

From the previous proposition, we easily deduce the following.

Corollary 4.6. *If $\bigcap \{S_\lambda \mid \lambda \in \Lambda\}$ is a locally finite defining family of overrings (respectively, \mathfrak{t} -flat overrings) of an integral domain D , and if each of S_λ is a Mori (respectively, strong Mori) domain, then D is a Mori (respectively, strong Mori) domain.*

Remark 4.7. Note that the Mori domain case in Corollary 4.6 can be viewed as a “non-completely integrally closed version” of the following well-known result [18, Proposition 1.4]: If $\bigcap \{S_\lambda \mid \lambda \in \Lambda\}$ is a locally finite defining family of an integral domain D and if each of S_λ a Krull domain, then D is a Krull domain.

Moreover, recall that Dessagnes in 1987 proved that the intersection of any locally finite family of Mori domains, all contained in the same integral domain, is a Mori domain [6, Proposition 3.2] (see also [35, Théorème 1]).

Recall that an integral domain D is a *weakly Krull domain* if $D = \bigcap \{D_P \mid P \in X^1(D)\}$, where $X^1(D)$ denotes the set of height one primes of D , and the intersection is locally finite. Weakly Krull domains were studied in [4].

It is well known that if D is a Mori domain then so is each of its rings of fractions [35, Théorème 2]. Using this piece of information and Proposition 4.5, we deduce immediately the following.

Corollary 4.8. *A weakly Krull domain D is a Mori domain if and only if D_P is a Mori domain for each $P \in X^1(D)$.*

5. Essential Domains and Prüfer v -multiplication Domains

In this section, we introduce a weak form of the finite character property of a defining family of a domain. As an application, we shed new light on the question of when an essential domain is a PvMD solved recently by Finocchiaro and Tartarone [9] using topological methods.

Let D be an integral domain, let $\mathcal{E}(D) := \{P \in \text{Spec}(D) \mid D_P \text{ a valuation domain}\}$ be the set of all essential valuation overrings of D , and let $\emptyset \neq X \subseteq \text{Spec}(D)$. We say that the domain D is X -essential if $X \subseteq \mathcal{E}(D)$ and $D = \bigcap \{D_P \mid P \in X\}$.

Recall from [5] that a prime ideal Q of D is an *associated prime of a principal ideal aD of D* , if Q is minimal over $(aD : bD)$ for some $b \in D \setminus aD$. For brevity, we call Q an *associated prime of D* and we denote by $\text{Assp}(D)$ the set of the associated prime ideals of D . We say that D is a P -domain if, for every $Q \in \text{Assp}(D)$, D_Q is a valuation domain [33]. Note that a PvMD is a P -domain and not conversely [33, Corollary 1.4, Example 2.1].

As we remarked above an important class of classical domains are X -essential for some nonempty set $X \subseteq \text{Spec}(D)$, i.e. weakly Krull domains, for $X = X^1(D)$. Moreover, if $X = \text{Max}(D)$ (or, even, $X = \text{Spec}(D)$) (respectively, $X = \text{Max}^f(D)$; $X = \text{Assp}(D)$) we get Prüfer domains (respectively, PvMDs; P -domains).

Let D be an X -essential domain, the (semi)star operation on D , $*_X$, induced by the nonempty family of overrings $\mathcal{X} := \{D_P \mid P \in X\}$, i.e. $*_X := \wedge_{\mathcal{X}}$ (defined by $E^{*_X} := \bigcap \{ED_P \mid P \in X\}$ for each $E \in \overline{\mathbf{F}}(D)$), is crucial for studying these domains as the following proposition shows.

Proposition 5.1. *Let D be an integral domain, let $\emptyset \neq X \subseteq \text{Spec}(D)$ such that $D = \bigcap \{D_P \mid P \in X\}$ and $*_X$ the star operation on D induced by the family of overrings $\{D_P \mid P \in X\}$. Then, the following are equivalent.*

- (i) D is an X -essential domain.
- (ii) Every $*_X$ -finite ideal is $*_X$ -invertible.

Proof. (i) \Rightarrow (ii) Let $I \in \mathbf{f}(D)$ and $P \in X$. Then $II^{-1}D_P = ID_P(ID_P)^{-1} = D_P$ since D_P is a valuation domain. Hence, $(II^{-1})^{*_X} = D$.

(ii) \Rightarrow (i) Let $P \in X$ and J a nonzero finitely generated ideal of D_P . Then $J = ID_P$ for some finitely generated ideal I of D . We have $JJ^{-1} = ID_P(ID_P)^{-1} = (II^{-1})D_P = (II^{-1})^{*_X}D_P = D_P$. So D_P is a local Prüfer domain, and hence a valuation domain. \square

S. El Baghdadi, M. Fontana & M. Zafrullah

Remark 5.2. Note that Proposition 5.1 provides a general setting for a well-known result on Prüfer domains (i.e. for $X = \text{Max}(D)$ and $*_X = \mathfrak{d}$) or on PvMDs (i.e. for $X = \text{Max}^t(D)$ and $*_X = \mathfrak{w}$). On the other hand, a Dedekind domain (respectively, a Krull domain) is a Prüfer Noetherian domain (respectively, a PvMD Mori (\mathfrak{w} -Noetherian) domain; note that in this situation $\mathfrak{t} = \mathfrak{w}$), that is, in both cases, X -essential and $*_X$ -Noetherian domain. We call a X -Dedekind domain an integral domain with these latter properties. Since on an X -essential and $*_X$ -Noetherian domain, the (semi)star operation $*_X$ is of finite type, then $*_X \leq \mathfrak{t}$ and so $\text{Max}^t(D) \subseteq \text{Max}^{*_X}(D)$ [4, Lemma 2.1]. Hence an X -Dedekind domain D is always a PvMD with $*_X = \mathfrak{w}$. Thus, Dedekind domains (i.e. when $*_X = \mathfrak{w} = \mathfrak{d}$) and Krull domains (i.e. when $*_X = \mathfrak{w}$) are the only X -Dedekind domains.

We say that a defining family $\{S_\lambda \mid \lambda \in \Lambda\}$ of an integral domain D has *GV-finite character property* if, for each ideal I of D such that $IS_\lambda = S_\lambda$ for every $\lambda \in \Lambda$, there exists a finitely generated ideal $J \subseteq I$ of D such that $JS_\lambda = S_\lambda$ for every λ . Note that the abbreviation “GV” stands for Glaz–Vasconcelos, since we will see that the GV-finite character property can be characterized by a general version of the notion of H -domain, introduced by Glaz and Vasconcelos in [22].

Obviously, every defining family of overrings of a Noetherian domain has GV-finite character property. Note that GV-finite character property is an extension of the finite character property. Indeed, assume that $D = \bigcap \{S_\lambda \mid \lambda \in \Lambda\}$ has the finite character property and let I be an ideal of D such that $IS_\lambda = S_\lambda$ for every λ . Let $0 \neq x \in I$ and let $\{S_{\lambda_k} \mid 1 \leq k \leq n\}$ be the only λ 's such that $xS_\lambda \neq S_\lambda$. For each k , there exists $J_k \subseteq I$ a finitely generated ideal of D such that $IS_{\lambda_k} = J_k S_{\lambda_k} = S_{\lambda_k}$. Take J to be the finitely generated subideal of I generated by x and the J_k 's, for $1 \leq k \leq n$, then it is straightforward that $JS_\lambda = IS_\lambda$ for each $\lambda \in \Lambda$.

Proposition 5.3. *Let D be an integral domain and let $\mathfrak{S} := \{S_\lambda \mid \lambda \in \Lambda\}$ be a defining family of overrings of D . Denote by $*$ the (semi)star operation on D induced by the defining family of overrings \mathfrak{S} of D , i.e. $*$:= $\wedge_{\mathfrak{S}}$. Then, the following are equivalent.*

- (i) \mathfrak{S} has GV-finite character property;
- (ii) for every ideal I of D such that $I^* = D$, there exists a finitely generated J ideal of D such that $J \subseteq I$ and $J^* = D$;
- (iii) the stable (semi)star operation $\bar{*}$, canonically associated to $*$, is of finite type, i.e. $\bar{*} = \tilde{*}$.

Proof. (i) \Leftrightarrow (ii) is straightforward.

(ii) \Rightarrow (iii) Let $E \in \overline{\mathfrak{F}}(D)$ and $x \in E^*$. Let I be a nonzero ideal of D such that $xI \subseteq E$ with $I^* = D$. By assumption, we can take I finitely generated. Let $F := xI \in \mathfrak{f}(D)$. Then $F \subseteq E$ and $x \in F^*$. Thus $\bar{*}$ is of finite type.

(iii) \Rightarrow (ii) is an easy consequence of the definitions. □

The case when D has a defining family of quotient rings, that is $D = \bigcap \{D_P \mid P \in X\}$ for some $X \subseteq \text{Spec}(D)$ is of particular interest. In this case, if the defining family $\{D_P \mid P \in X\}$ of D has GV-finite character property, we simply say that the subset X of $\text{Spec}(D)$ has *GV-finite character property*. Note that, in this case, $*$ is necessarily stable, that is $\bar{*} = *$. Clearly, for any domain D , the sets $\text{Max}(D)$ and $\text{Max}^t(D)$ have GV-finite character property. Therefore, from Proposition 5.3 and from [8, Corollary 2.8, Proposition 2.9], we easily deduce the following.

Corollary 5.4. *Let D be an integral domain and let $\mathfrak{X} := \{D_P \mid P \in X\}$ be a defining family of quotient rings of D for some nonempty $X \subseteq \text{Spec}(D)$. Let $*_X$ be the (semi)star operation on D , induced by the family of overrings $\mathfrak{X} := \{D_P \mid P \in X\}$, i.e. $*_X := \wedge_{\mathfrak{X}}$. Then the following are equivalent.*

- (i) X has GV-finite character property;
- (ii) If I is an ideal of D such that $I \not\subseteq P$ for every ideal $P \in X$, then there exists $J \subseteq I$ a finitely generated ideal of D such that $J \not\subseteq P$ for every ideal $P \in X$;
- (iii) $*_X$ is of finite type;
- (iv) X is quasi-compact for the Zariski topology on $\text{Spec}(R)$.

Given a semistar operation \star on an integral domain D , D is called an $H(\star)$ -domain [17] if for every nonzero ideal I of D such that $I^\star = D$, there exists a nonzero finitely generated ideal J of D such that $J \subseteq I$ and $J^\star = D$. Thus, given an integral domain D and $X \subseteq \text{Spec}(D)$ such that $\{D_P \mid P \in X\}$ is a defining family of quotient rings of D , by Proposition 5.3, X has GV-finite character property if and only if D is an $H(*_X)$ -domain, where $*_X$ is the (semi)star operation induced by the defining family $\{D_P \mid P \in X\}$ of D .

Note that the $H(\star)$ -domains generalize in the semistar setting the H -domains introduced by Glaz and Vasconcelos [22]; more precisely, the H -domains coincide with the $H(\mathfrak{v})$ -domains [17, Sec. 2]

The following theorem provides an algebraic version of the solution of the problem when an essential domain is a PvMD. This problem was recently solved in [9] using topological methods.

Theorem 5.5. *Let D be an integral. Then the following are equivalent.*

- (i) D is a PvMD;
- (ii) D is essential and the set $\{D_P \mid P \in \mathcal{E}(D)\}$ of all essential valuation overrings of D has GV-finite character property.
- (iii) D is essential and, for all $a, b \in D \setminus \{0\}$, $aD \cap bD = F^{\mathfrak{v}}$ for some $F \in \mathfrak{f}(D)$ (in particular, $F \subseteq aD \cap bD$).

Proof. (i) \Rightarrow (ii) Since a PvMD is an essential domain, we next show that $\mathcal{E}(D)$ has GV-finite character property. Let I be an ideal of D such that $I \not\subseteq P$ for every $P \in \text{Spec}(D)$ such that $D_P \in \mathcal{E}(D)$ (such a prime ideal is called *essential prime* of D). Since $\text{Max}^t(D) \subseteq \mathcal{E}(D)$, $*_{\mathcal{E}(D)} \leq *_{\text{Max}^t(D)} = \mathfrak{w}$. Hence $I^* = D$. Then, there

S. El Baghdadi, M. Fontana & M. Zafrullah

exists a nonzero finitely generated ideal J of D such that $J \subseteq I$ and $J^* = D$. But, as each essential prime ideal P is such that PD_P is a \mathfrak{t} -ideal in the valuation domain D_P , P is a \mathfrak{t} -ideal of D [29, Lemma 3.17] and so it is contained in a maximal \mathfrak{t} -ideal. Thus, we get that $J \not\subseteq P$ for every essential prime P . Therefore, $\mathcal{E}(D)$ has GV-finite character property.

(ii) \Rightarrow (i) By assumption, we have $D = \bigcap \{D_P \mid P \in \mathcal{E}(D)\}$. By Corollary 5.4, the (semi)star operation $*_{\mathcal{E}(D)}$ is of finite type, so $*_{\mathcal{E}(D)} \leq \mathfrak{t}$. Hence, each \mathfrak{t} -maximal ideal is a $*_{\mathcal{E}(D)}$ -ideal. Thus, each \mathfrak{t} -maximal ideal is contained in an essential prime ideal, and hence it is an essential prime. This proves that D is a PvMD.

(i) \Rightarrow (iii) Recall that $aD \cap bD = ab(a, b)^{-1}$. Since D is a PvMD, we have $((a, b)(a, b)^{-1})^* = D$. By a standard argument, we can find a finitely generated subideal F of $aD \cap bD$ such that $aD \cap bD = F^* = F^v$.

(iii) \Rightarrow (i) is well known [39, Lemma 8]. \square

Remark 5.6. By the above characterization, an essential domain to be a PvMD is equivalent to the condition that the (semi)star operation induced by the defining family is of finite type, and in this case it is the \mathfrak{w} -operation.

A P-domain need not be a PvMD, see an example in [33]. This shows that the defining family of localizations at associated primes of a P-domain does not have in general GV-finite character property, or equivalently, the (semi)star operation induced by this defining family is not in general of finite type.

Acknowledgments

The authors thank the referee for the evaluation of the paper and for the helpful suggestions. The second author thanks the “National Group for Algebraic and Geometric Structures, and their Applications” (GNSAGA) of the *Istituto Nazionale di Alta Matematica* for a partial support.

References

- [1] D. D. Anderson, Star operations induced by overrings, *Comm. Algebra* **16** (1988) 2535–2553.
- [2] D. D. Anderson and D. F. Anderson, Some remarks on star operations and the class group, *J. Pure Appl. Algebra* **51** (1988) 27–33.
- [3] D. D. Anderson and S. J. Cook, Two star-operations and their induced lattices, *Comm. Algebra* **28** (2000) 2461–2475.
- [4] D. D. Anderson, J. Mott and M. Zafrullah, Finite character representations for integral domains, *Boll. Unione Mat. Ital. (7)* **6-B** (1992) 613–630.
- [5] J. Brewer and W. Heinzer, Associated primes of principal ideals, *Duke Math. J.* **41** (1974) 1–7.
- [6] N. Dessagnes, Intersections d’anneaux de Mori-exemples, *Port. Math.* **44** (1987) 379–392.
- [7] S. El Baghdadi, M. Fontana and G. Picozza, Semistar Dedekind domains. *J. Pure Appl. Algebra* **193** (2004) 27–60.
- [8] C. A. Finocchiaro and D. Spirito, Some topological considerations on semistar operations, *J. Algebra* **409** (2014) 199–218.

- [9] C. A. Finocchiaro and F. Tartarone, On a topological characterization of Prüfer v -multiplication domains among essential domains, preprint (2014), arXiv:1410.4037.
- [10] M. Fontana and S. Gabelli, On the class group and the local class group of a pullback, *J. Algebra* **181** (1996) 803–835.
- [11] M. Fontana and J. Huckaba, Localizing systems and semistar operations, in *Non-Noetherian Commutative Ring Theory*, Mathematics and its Applications, Vol. 520 (Kluwer Academic Publishers Dordrecht, 2000), pp. 169–197.
- [12] M. Fontana, P. Jara and E. Santos, Prüfer \star -multiplication domains and semistar operations, *J. Algebra Appl.* **2** (2003) 21–50.
- [13] M. Fontana and K. A. Loper, Kronecker function rings: A general approach, in *Ideal Theoretic Methods in Commutative Algebra*, Lecture Notes in Pure and Applied Mathematics, Vol. 220 (Marcel, Dekker, New York, 2001), pp. 189–205.
- [14] M. Fontana and K. A. Loper, N -rings, Kronecker function rings and related semistar operations, *Comm. Algebra* **31** (2003) 4775–4805.
- [15] M. Fontana and K. A. Loper, Cancellation properties in ideal systems: A classification of e.a.b. semistar operations, *J. Pure Appl. Algebra* **213** (2009) 2095–2103.
- [16] M. Fontana, K. A. Loper and R. Matsuda, Cancellation properties in ideal systems: An e.a.b. not a.b. star operation, *Arab. J. Sci. Eng.* **35** (2010) 45–49.
- [17] M. Fontana and G. Picozza, Semistar invertibility on integral domains, *Algebra Colloq.* **12** (2005) 645–664.
- [18] R. Fossum, *The Divisor Class Group of a Krull Domain*, Ergebnisse der Mathematik und ihrer Grenzgebiete, Vol. 74 (Springer-Verlag, Berlin, 1973).
- [19] J. M. García, P. Jara and E. Santos, Prüfer \ast -multiplication domains and torsion theories, *Comm. Algebra* **27** (1999) 1275–1295.
- [20] R. Gilmer, *Multiplicative Ideal Theory* (Marcel, Dekker, New York, 1972).
- [21] R. Gilmer and W. Heinzer, Intersections of quotient rings of an integral domain, *J. Math. Kyoto Univ.* **7** (1967) 133–150.
- [22] S. Glaz and W. V. Vasconcelos, Flat ideals. II, *Manuscripta Math.* **22** (1977) 325–341.
- [23] M. Griffin, Some results on v -multiplication rings, *Canad. J. Math.* **19** (1967) 710–722.
- [24] F. Halter-Koch, *Ideal Systems: An Introduction to Multiplicative Ideal Theory* (Marcel Dekker, New York, 1998).
- [25] W. Heinzer and J. Ohm, An essential ring which is not a v -multiplication ring, *Canad. J. Math.* **25** (1973) 856–861.
- [26] W. Heinzer and M. Roitman, Well-centered overrings of an integral domain, *J. Algebra* **272** (2004) 435–455.
- [27] E. Houston, S. Malik and J. Mott, Characterizations of \ast -multiplication domains, *Canad. Math. Bull.* **27** (1984) 48–52.
- [28] P. Jaffard, *Les Systèmes d’Idéaux* (Dunod, Paris, 1960).
- [29] B. G. Kang, Prüfer v -Multiplication domains and the ring $R[X]_{N_v}$, *J. Algebra* **123** (1989) 151–170.
- [30] D. J. Kwak and Y. S. Park, On t -flat overrings, *Chinese J. Math.* **23**(1) (1995) 17–24.
- [31] S. Malik, J. Mott and M. Zafrullah, On t -invertibility, *Comm. Algebra* **16** (1988) 149–170.
- [32] H. Matsumura, *Commutative Ring Theory* (Cambridge University Press, Cambridge, 1986).
- [33] J. Mott and M. Zafrullah, On Prüfer v -multiplication domains, *Manuscripta Math.* **35** (1981) 1–26.
- [34] G. Picozza, Semistar operations and multiplicative ideal theory, Ph.D. Thesis, Università degli Studi “Roma Tre”, Rome (2004).

S. El Baghdadi, M. Fontana & M. Zafrullah

- [35] J. Querré, Intersections d'anneaux intègres, *J. Algebra* **43** (1976) 55–60.
- [36] F. Richman, Generalized quotient rings, *Proc. Amer. Math. Soc.* **16** (1965) 794–799.
- [37] H. Uda, LCM-stableness in ring extensions, *Hiroshima Math. J.* **13** (1983) 357–377.
- [38] F. Wang and R. McCasland, On w -modules over strong Mori domains, *Comm. Algebra* **25** (1997) 1285–1306.
- [39] M. Zafrullah, On finite conductor domains, *Manuscripta Math.* **24** (1978) 191–203.
- [40] M. Zafrullah, Ascending chain conditions and star operations, *Comm. Algebra* **17** (1989) 1523–1533.
- [41] M. Zafrullah, Putting t -invertibility to use, in *Non-Noetherian Commutative Ring Theory*, Mathematics and its Applications, Vol. 520 (Kluwer Academic Publishers Dordrecht, 2000), pp. 429–457.