

SPECTRAL SPACES OF SEMISTAR OPERATIONS

CARMELO A. FINOCCHIARO, MARCO FONTANA, AND DARIO SPIRITO

ABSTRACT. We investigate, from a topological point of view, the classes of spectral semistar operations and of **eab** semistar operations, following methods recently introduced in [11, 13]. We show that, in both cases, the subspaces of finite type operations are spectral spaces in the sense of Hochster and, moreover, that there is a distinguished class of overrings strictly connected to each of the two types of collections of semistar operations. We also prove that the space of stable semistar operations is homeomorphic to the space of Gabriel-Popescu localizing systems, endowed with a Zariski-like topology, extending to the topological level a result established in [14]. As a side effect, we obtain that the space of localizing systems of finite type is also a spectral space. Finally, we show that the Zariski topology on the set of semistar operations is the same as the b -topology defined recently by B. Olberding [37, 38].

1. INTRODUCTION

In 1936, W. Krull introduced, in his first *Beiträge* paper [32] (see also [33]), the concept of a “special” closure operation on the nonzero fractional ideals, called star operation. If D is an integral domain with quotient field K , in 1994, Okabe and Matsuda [35] suggested the terminology of semistar operation for a more “flexible” and general notion of a closure operation \star , defined on the set of nonzero D -submodules of K , allowing $D \neq D^\star$. However, it is worth noting that this kind of operation was previously considered by J. Huckaba, in the very general setting of rings with zero divisors [25, Section 20] (cf. also [23, Section 32], [1], [2], [3], [9], [10], [16], [26], [27], [28], [29]).

The set of semistar operations on a domain D can be endowed with a topology (called the *Zariski topology*), as in [13], in such a way that both the prime spectrum of D and the set of overrings of D are naturally topologically embedded in it. This topology was used to study the problem of when the semistar operation defined by a family of overrings is of finite type [5, Problem 44]. Subsequently, it was proved that the set of semistar operations of finite type is a spectral space.

The purpose of this paper is to deepen and specialize the study of the Zariski topology on $\mathbf{SStar}(D)$ to the case of the distinguished subspaces $\overline{\mathbf{SStar}}(D)$ (Section 3 and 4) and $\mathbf{SStar}_{\text{eab}}(D)$ (Section 5) comprising, respectively, the stable and the **eab** semistar operations. We will show that, in both cases, there is a topological retraction to the set of finite type operations and that there is a distinguished class of overrings connected to each of the two types of collections of semistar operations. We will also show that both the set of finite type spectral operations and the set of

2010 *Mathematics Subject Classification.* 13A15, 13G05, 13B10, 13E99, 13C11, 14A05.

Key words and phrases. Spectral spaces, Riemann-Zariski space of valuation domains, semistar operations, Zariski topology, inverse topology, ultrafilter topology, localizing system.

This work was partially supported by GNSAGA of *Istituto Nazionale di Alta Matematica*.

finite type **eab** operations are spectral spaces, reducing the latter case to the former. However, the proofs given here are not constructive, and provide only vague hints on how such a ring might look like. We also prove that the space of stable semistar operations is homeomorphic to the space of Gabriel-Popescu localizing systems, endowed with a natural topology (described later), extending to the topological level a result established in [14]. As a side effect, we obtain that the space of localizing systems of finite type is also a spectral space. Finally, we show that the Zariski topology on the set of semistar operations is the same as the b -topology defined by Olberding [37, 38].

2. PRELIMINARIES

Throughout this paper, let D be an integral domain with quotient field K . Let $\overline{\mathbf{F}}(D)$ [respectively, $\mathbf{F}(D)$; $\mathbf{f}(D)$] be the set of all nonzero D -submodules of K [respectively, nonzero fractional ideals; nonzero finitely generated fractional ideals] of D (thus, $\mathbf{f}(D) \subseteq \mathbf{F}(D) \subseteq \overline{\mathbf{F}}(D)$).

A mapping $\star : \overline{\mathbf{F}}(D) \rightarrow \overline{\mathbf{F}}(D)$, $E \mapsto E^\star$, is called a *semistar operation* of D if, for all $z \in K$, $z \neq 0$ and for all $E, F \in \overline{\mathbf{F}}(D)$, the following properties hold: (\star_1) $(zE)^\star = zE^\star$; (\star_2) $E \subseteq F \Rightarrow E^\star \subseteq F^\star$; (\star_3) $E \subseteq E^\star$; and (\star_4) $E^{\star\star} := (E^\star)^\star = E^\star$.

When $D^\star = D$, the restriction of \star to $\mathbf{F}(D)$ is called a *star operation* (see [23, Section 32] for more details).

As in the classical star-operation setting, we associate to a semistar operation \star of D a new semistar operation \star_f of D defining, for every $E \in \overline{\mathbf{F}}(D)$,

$$E^{\star_f} := \bigcup \{F^\star \mid F \subseteq E, F \in \mathbf{f}(D)\}.$$

We call \star_f the semistar operation of finite type of D *associated* to \star . If $\star = \star_f$, we say that \star is a semistar operation of *finite type* on D . Note that $(\star_f)_f = \star_f$, so \star_f is a semistar operation of finite type on D .

We denote by $\mathbf{SStar}(D)$ [respectively, $\mathbf{SStar}_f(D)$] the set of all semistar operations [respectively, semistar operations of finite type] on D . Given two semistar operations \star' and \star'' of D , we say that $\star' \preceq \star''$ if $E^{\star'} \subseteq E^{\star''}$, for all $E \in \overline{\mathbf{F}}(D)$. The relation “ \preceq ” introduces a partial ordering in $\mathbf{SStar}(D)$. From the definition of \star_f , we deduce that $\star_f \preceq \star$ and that \star_f is the largest semistar operation of finite type smaller or equal to \star .

Let \mathcal{S} be a nonempty set of semistar operations on D . For each $E \in \overline{\mathbf{F}}(D)$, define $E^{\wedge_{\mathcal{S}}}$ as follows:

$$E^{\wedge_{\mathcal{S}}} := \bigcap \{E^\star \mid \star \in \mathcal{S}\}.$$

It is easy to see that $E^{\wedge_{\mathcal{S}}}$ is a semistar operation on D and it is the infimum of \mathcal{S} in the partially ordered set $(\mathbf{SStar}(D), \preceq)$. The semistar operation $E^{\vee_{\mathcal{S}}} := \bigwedge \{\sigma \in \mathbf{SStar}(D) \mid \star \preceq \sigma \text{ for all } \star \in \mathcal{S}\}$ is the supremum of \mathcal{S} in $(\mathbf{SStar}(D), \preceq)$.

A nonzero ideal I of D is called a *quasi- \star -ideal* if $I = I^\star \cap D$. A *quasi- \star -prime* is a quasi- \star -ideal which is also a prime ideal. The set of all quasi- \star -prime ideals of D is denoted by $\mathbf{QSpec}^\star(D)$. The set of maximal elements in the set of proper quasi- \star -ideals of D (ordered by set-theoretic inclusion) is denoted by $\mathbf{QMax}^\star(D)$ and it is a subset of $\mathbf{QSpec}^\star(D)$. It is well known that if \star is a semistar operation of finite type then $\mathbf{QMax}^\star(D)$ is nonempty [16, Lemma 2.3(1)]. A semistar operation \star is called *quasi-spectral* (or *semifinite*) if each quasi- \star -ideal is contained in a quasi- \star -prime.

In [13], the set $\mathbf{SStar}(D)$ of all semistar operation was endowed with a topology (called the *Zariski topology*) having, as a subbasis of open sets, the sets of the type $\mathbf{V}_E := \{\star \in \mathbf{SStar}(D) \mid 1 \in E^\star\}$, where E is a nonzero D -submodule of K . This topology makes $\mathbf{SStar}(D)$ into a quasi-compact T_0 space.

For each overring T of D , we can define a semistar operation of finite type $\wedge_{\{T\}} : \overline{\mathbf{F}}(D) \rightarrow \overline{\mathbf{F}}(D)$ by setting $E^{\wedge_{\{T\}}} := ET$, for each $E \in \overline{\mathbf{F}}(D)$. If we have a whole family \mathcal{T} of overrings of D , we can consider the semistar operation $\wedge_{\mathcal{T}} := \wedge\{\wedge_{\{T\}} \mid T \in \mathcal{T}\}$.

For future reference, we state an embedding property that will be used later several times.

Proposition 2.1. *Let $\mathbf{Overr}(D)$ be the set of all overrings of D , endowed with the topology whose basic open sets are of the form $\mathbf{Overr}(D[x_1, x_2, \dots, x_n])$, for x_1, x_2, \dots, x_n varying in K .*

- (1) *The map $\iota : \mathbf{Overr}(D) \rightarrow \mathbf{SStar}_f(D)$, defined by $\iota(T) := \wedge_{\{T\}}$, for each $T \in \mathbf{Overr}(D)$, is a topological embedding [13, Proposition 2.5].*
- (2) *The map $\pi : \mathbf{SStar}_f(D) \rightarrow \mathbf{Overr}(D)$, defined by $\pi(\star) := D^\star$, for any $\star \in \mathbf{SStar}_f(D)$, is a continuous surjection.*
- (3) *$\pi \circ \iota$ is the identity map of $\mathbf{Overr}(D)$, that is, π is a topological retraction.*

Proof. Part (1) is [13, Proposition 2.5], while parts (2) and (3) follow from the fact that

$$\pi^{-1}(\mathbf{Overr}(D[x])) = \mathbf{SStar}_f(D) \cap \mathbf{V}_{x^{-1}D}. \quad \square$$

Remark 2.2. Note that, by [13, Proposition 2.4(2)], a statement completely analogous to Proposition 2.1 holds when $\mathbf{SStar}(D)$ replaces everywhere $\mathbf{SStar}_f(D)$.

Recall that two different points x, y of a topological space are *topologically distinguishable* if there is an open set which contains one of these points and not the other. Obviously, the previous property holds for any pair of distinct points if and only if the space is T_0 . On the other hand, “topological indistinguishability” of points is an equivalence relation. No matter what topological space X might be to begin with, the quotient space under this equivalence relation is always T_0 . This quotient space is called *the Kolmogoroff quotient space of X* .

Let X be a spectral space (i.e., a topological space that is homeomorphic to the prime spectrum of a ring, endowed with the Zariski topology). It is possible to consider on X another topology (see [24, Proposition 8]) defined by taking the collection of all the open and quasi-compact subspaces of X as a basis of closed sets. This topology is called *the inverse topology on X* . Note that, by definition, the closure of a subset Y of X , with respect to the inverse topology, is given by

$$\bigcap \{U \mid U \subseteq X \text{ open and quasi-compact, } Y \subseteq U\}.$$

3. STABLE SEMISTAR OPERATIONS AND LOCALIZING SYSTEMS

A semistar operation \star defined on an integral domain D is called *stable* provided that, for any $E, H \in \overline{\mathbf{F}}(D)$, we have $(E \cap H)^\star = E^\star \cap H^\star$. We denote by $\overline{\mathbf{SStar}}(D)$ the set of stable semistar operations on D .

Proposition 3.1. *Let D be an integral domain and let $\iota : \mathbf{Overr}(D) \rightarrow \mathbf{SStar}_f(D)$ be the topological embedding defined in Proposition 2.1. If $T \in \mathbf{Overr}(D)$, then $\iota(T) \in \overline{\mathbf{SStar}}(D)$ if and only if T is flat over D .*

Proof. It is enough to note that the equality $(I \cap J)T = IT \cap JT$ holds for every ideal I, J of D if and only if T is flat ([40, Proposition 1.7] and [34, Theorem 7.4(i)]). \square

Given a semistar operation \star on D , we can always associate to \star a stable semistar operation $\overline{\star}$ by defining, for every $E \in \overline{\mathbf{F}}(D)$,

$$E^{\overline{\star}} := \bigcup \{(E : I) \mid I \text{ nonzero ideal of } D \text{ such that } I^{\star} = D^{\star}\}.$$

It is easy to see that $\overline{\star} \preceq \star$ and, moreover, that $\overline{\star}$ is the largest stable semistar operation that precedes \star . Therefore, \star is stable if and only if $\star = \overline{\star}$ [14, Proposition 3.7, Corollary 3.9].

Proposition 3.2. *Let D be an integral domain, and denote by \mathcal{I} the set of nonzero ideals of D . Let also $\overline{\Phi} : \mathbf{SStar}(D) \rightarrow \overline{\mathbf{SStar}}(D)$ be the map defined by $\overline{\Phi}(\star) := \overline{\star}$, for each $\star \in \mathbf{SStar}(D)$.*

- (1) *The set $\{\mathbf{V}_I \cap \overline{\mathbf{SStar}}(D) \mid I \in \mathcal{I}\}$ is a subbasis for $\overline{\mathbf{SStar}}(D)$.*
- (2) *If $\mathbf{SStar}(D)$ is endowed with the Zariski topology, $\overline{\Phi}$ is a topological retraction.*
- (3) *If $\mathbf{SStar}(D)$ is endowed with the topology generated by the family $\{\mathbf{V}_I \mid I \in \mathcal{I}\}$, then $\overline{\Phi}$ is the canonical map onto the Kolmogoroff quotient space of $\mathbf{SStar}(D)$.*

Proof. (1) For any nonzero D -submodule E of K , and any stable semistar operation \star , we have $1 \in E^{\star}$ if and only if $1 \in E^{\star} \cap D^{\star} = (E \cap D)^{\star}$. Therefore, $\star \in \mathbf{V}_E$ if and only if $\star \in \mathbf{V}_{E \cap D}$. The claim follows.

(2) We claim that, if I is an ideal of D , $\overline{\Phi}^{-1}(\mathbf{V}_I \cap \overline{\mathbf{SStar}}(D)) = \mathbf{V}_I$. Indeed, if $\star \in \mathbf{V}_I$ then $1 \in I^{\star}$, so $1 \in (I : I) \subseteq I^{\overline{\star}}$ and $\overline{\Phi}(\star) \in \mathbf{V}_I \cap \overline{\mathbf{SStar}}(D)$. Conversely, if $\star \in \overline{\Phi}^{-1}(\mathbf{V}_I \cap \overline{\mathbf{SStar}}(D))$, then $\overline{\Phi}(\star) \in \mathbf{V}_I$, and thus $1 \in (I : E)$ for some $E \in \mathcal{I}$ such that $E^{\star} = D^{\star}$. But this means that $E \subseteq I$, and so $1 \in I^{\star}$, i.e., $\star \in \mathbf{V}_I$. Hence, $\overline{\Phi}$ is continuous.

Since \star is stable if and only if $\star = \overline{\star}$ [14, Proposition 3.7, Corollary 3.9], it follows that $\overline{\Phi}$ is a topological retraction.

(3) Since, by the previous point, $\overline{\Phi}^{-1}(\mathbf{V}_I \cap \overline{\mathbf{SStar}}(D)) = \mathbf{V}_I$, the map $\overline{\Phi}$ is continuous even when $\mathbf{SStar}(D)$ is endowed with the weaker topology. To show that $\overline{\Phi}$ is the canonical map onto the Kolmogoroff quotient space of $\mathbf{SStar}(D)$, it is enough to show that $\overline{\star}_1 = \overline{\star}_2$ if and only if $\star_1 \in \mathbf{V}_I$ is equivalent to $\star_2 \in \mathbf{V}_I$.

Suppose $\overline{\star}_1 = \overline{\star}_2$, and let I be an ideal of D such that $\star_1 \in \mathbf{V}_I$, that is, $1 \in I^{\star_1}$. By definition, $1 \in (I : I) \subseteq I^{\overline{\star}_1} = I^{\overline{\star}_2} \subseteq I^{\star_2}$. It follows that $\star_2 \in \mathbf{V}_I$. By symmetry, we deduce that if $\star_2 \in \mathbf{V}_I$ then $\star_1 \in \mathbf{V}_I$.

Conversely, suppose that $\star_1 \in \mathbf{V}_I$ if and only if $\star_2 \in \mathbf{V}_I$. Then, $I^{\star_1} = D^{\star_1}$ if and only if $I^{\star_2} = D^{\star_2}$; a direct application of the definition of the stable semistar operations canonically associated shows that $\overline{\star}_1 = \overline{\star}_2$ (cf. [14, page 182]). \square

Remark 3.3. Recall that for any *star* operation \star on an integral domain D with quotient field K and for any $E \in \overline{\mathbf{F}}(D)$, we can consider the map \star_e defined by

$$E^{\star_e} := \begin{cases} E^{\star} & \text{if } E \in \mathbf{F}(D), \\ K & \text{if } E \in \overline{\mathbf{F}}(D) \setminus \mathbf{F}(D). \end{cases}$$

The map $E \mapsto E^{\star_e}$ defines a semistar operation on D such that $D^{\star_e} = D$, called *the trivial semistar extension of \star* . Note that, even if \star is a stable star operation,

$*_e$ is not always stable: for example, let D be a Dedekind domain with exactly two maximal ideals, P and Q , and let $*$ be the identity star operation. Then, D_P and D_Q are not fractional ideals of D [13, Example 5.7], and thus $D_P^{*e} = K = D_Q^{*e}$. On the other hand, $(D_P \cap D_Q)^{*e} = D^{*e} = D$.

Proposition 3.4. *Let $*$ be a stable star operation on an integral domain D . There is exactly one stable semistar operation \star on D such that $\star|_{\mathbf{F}(D)} = *$.*

Proof. Suppose there exist two stable semistar extensions \star_1 and \star_2 of the star operation $*$, i.e., $\star_1|_{\mathbf{F}(D)} = \star_2|_{\mathbf{F}(D)} = *$. Since $\overline{\mathbf{SStar}}(D)$ is T_0 , there is a subbasic open set $U_I := V_I \cap \overline{\mathbf{SStar}}(D)$, with I a proper ideal of D , such that $\star_1 \in U_I$ but $\star_2 \notin U_I$ (or conversely). But this would imply $I^* = I^{\star_1} \neq I^{\star_2} = I^*$, which is absurd.

For the existence, consider the semistar operation $\star := \overline{*_e}$, where $*_e$ is the trivial semistar extension of $*$ defined in Remark 3.3; by definition, \star is a stable semistar operation. On the other hand, since $D^* = D$, if I is a nonzero D -submodule of K such that $I^* = D^*$, then I is an ideal in D . It follows that $\star|_{\mathbf{F}(D)}$ is the stable closure of $*$ as a *star* operation, as defined in [2, Definition 2.2]. However, since $*$ is already stable, we have $\star|_{\mathbf{F}(D)} = *$, i.e., \star is an extension of $*$. \square

Our next goal is to establish a topological connection between stable operations and localizing systems.

A *localizing system* on D is a subset \mathcal{F} of ideals of D such that:

- if $I \in \mathcal{F}$ and J is an ideal of D such that $I \subseteq J$, then $J \in \mathcal{F}$;
- if $I \in \mathcal{F}$ and J is an ideal of D such that, for each $i \in I$, $(J :_D iD) \in \mathcal{F}$, then $J \in \mathcal{F}$.

A localizing system \mathcal{F} is *of finite type* if for each $I \in \mathcal{F}$ there exists a nonzero finitely generated ideal $J \in \mathcal{F}$ with $J \subseteq I$. For instance, if T is an overring of R , $\mathcal{F}(T) := \{I \mid I \text{ ideal of } D, IT = T\}$ is a localizing system of finite type. On the other hand, if V is a valuation domain and P is a nonzero idempotent prime ideal of V , then $\hat{\mathcal{F}}(P) := \{I \mid I \text{ ideal of } V \text{ and } I \supseteq P\}$ is a localizing system of V which is not of finite type. Given a localizing system \mathcal{F} of an integral domain D , then

$$\mathcal{F}_f := \{I \in \mathcal{F} \mid I \supseteq J, \text{ for some nonzero finitely generated ideal } J \in \mathcal{F}\}$$

is a localizing system of finite type of D , and $\mathcal{F} = \mathcal{F}_f$ if and only if \mathcal{F} is a localizing system of finite type.

We denote by $\mathbf{LS}(D)$ [respectively, $\mathbf{LS}_f(D)$] the set of all localizing systems [respectively, localizing systems of finite type] on D . For further details on localizing systems, see [4, Chap. II, §2, Exercices 17–25] or [14, Sections 2 and 3].

It is well known that, to each localizing system \mathcal{F} , we can associate a semistar operation $\star_{\mathcal{F}}$ defined as follows, for each $E \in \overline{\mathbf{F}}(D)$,

$$E^{\star_{\mathcal{F}}} := \bigcup \{(E : H) \mid H \in \mathcal{F}\}.$$

The assignment $\mathcal{F} \mapsto \star_{\mathcal{F}}$ defines a map $\lambda : \mathbf{LS}(D) \rightarrow \mathbf{SStar}(D)$. By [14, Theorem 2.10, Corollary 2.11, and Proposition 3.2], λ is an injective map whose image is exactly $\overline{\mathbf{SStar}}(D)$.

On the set $\mathbf{LS}(D)$ of localizing system on D we can introduce a natural topology, that we still call the *Zariski topology*, whose subbasic open sets are the $W_I := \{\mathcal{F} \in \mathbf{LS}(D) \mid I \in \mathcal{F}\}$, as I varies among the ideals of D .

Proposition 3.5. *Let D be an integral domain. The map $\lambda : \text{LS}(D) \rightarrow \overline{\text{SStar}}(D)$, $\mathcal{F} \mapsto \star_{\mathcal{F}}$, establishes a homeomorphism between spaces endowed with the Zariski topologies.*

Proof. By the previous remarks, we only need to show that λ is continuous and open. Let U_I be a subbasic open set of $\overline{\text{SStar}}(D)$. Then,

$$\begin{aligned} \lambda^{-1}(U_I) &= \{\mathcal{F} \in \text{LS}(D) \mid \star_{\mathcal{F}} \in U_I\} = \\ &= \{\mathcal{F} \in \text{LS}(D) \mid 1 \in (I : H) \text{ for some } H \in \mathcal{F}\} = \\ &= \{\mathcal{F} \in \text{LS}(D) \mid H \subseteq I \text{ for some } H \in \mathcal{F}\} = \bigcup_{H \subseteq I} W_H, \end{aligned}$$

and thus λ is open. Moreover, $\bigcup_{H \subseteq I} W_H = W_I$: indeed, if $\mathcal{F} \in W_H$ then $H \in \mathcal{F}$, and so $I \in \mathcal{F}$, while the left hand union trivially contains W_I . Therefore, $\lambda^{-1}(U_I) = W_I$, and, since λ is bijective, $\lambda(W_I) = U_I$. Therefore, λ is both continuous and open, and thus a homeomorphism. \square

4. SPECTRAL SEMISTAR OPERATIONS

If Y is a subset of the prime spectrum $\text{Spec}(D)$ of an integral domain D , then we define the semistar operation s_Y induced by Y as the semistar operation associated to the set $\mathcal{T}(Y) := \{D_P \mid P \in Y\}$, i.e.,

$$E^{s_Y} := \bigcap \{ED_P \mid P \in Y\}, \text{ for every } E \in \overline{\mathbf{F}}(D).$$

If $Y = \emptyset$, we have an empty intersection, and we set as usual $E^{s_\emptyset} := K$ for every $E \in \overline{\mathbf{F}}(D)$ (or, equivalently, $s_\emptyset := \wedge_{\{K\}}$).

A semistar operation of the type s_Y , for some $Y \subseteq \text{Spec}(D)$, is called a *spectral semistar operation*.

Note that, if we take $Y = \{(0)\}$, we also have that $s_Y = \wedge_{\{K\}}$. Therefore, without loss of generality, in the definition of spectral semistar operation we can assume $\emptyset \neq Y \subseteq \text{Spec}(D)$.

Denote by $\text{SStar}_{sp}(D)$ the set of spectral semistar operations, and by $\text{SStar}_{f,sp}(D)$ the set of spectral semistar operations of finite type.

Since each localization of D is D -flat, and the infimum of a family of stable semistar operation is again stable, using Proposition 3.1 we see that every spectral semistar operation is stable. On the other hand, not every stable operation is spectral; however, a stable semistar operation is spectral if and only if it is quasi-spectral [14, Proposition 4.23(2)]. In particular, every finite type stable operation is spectral (cf. [1, Corollary 4.2] and [14, page 185 and Theorem 4.12(3)]), so that $\text{SStar}_{f,sp}(D)$ coincides with the set of stable operations of finite type (sometimes denoted by $\widetilde{\text{SStar}}(D)$).

Like for $\overline{\star}$, we can associate to each semistar operation \star a stable semistar operation of finite type $\widetilde{\star}$ by defining, for every $E \in \overline{\mathbf{F}}(D)$,

$$E^{\widetilde{\star}} := \bigcup \{(E : J) \mid J \text{ nonzero finitely generated ideal of } D \\ \text{such that } J^{\star} = D^{\star}\}.$$

The stable semistar operation of finite type $\widetilde{\star}$ is smaller than \star , and it is the biggest stable semistar operation of finite type smaller than \star . It follows that \star is stable of finite type if and only if $\star = \widetilde{\star}$.

In the following proposition, we collect some of the properties concerning the relation between Y and s_Y . As usual, for each subset Y of $\text{Spec}(D)$, we set $Y^{\text{gen}} :=$

$\{z \in \text{Spec}(D) \mid y \in \mathbf{Cl}(\{z\}) \text{ for some } y \in Y\}$ and we denote $\mathbf{Cl}^{\text{inv}}(Y)$ the closure of Y in the inverse topology of $\text{Spec}(D)$.

Proposition 4.1. (cf. [13, Corollaries 4.4 and 5.2, Proposition 5.1] and [14, Lemma 4.2 and Remark 4.5]) *Let D be an integral domain and let Y and Z be two nonempty subsets of $\text{Spec}(D)$. The following statements hold.*

- (1) $s_Y = s_Z$ if and only if $Y^{\text{gen}} = Z^{\text{gen}}$.
- (2) s_Y is of finite type if and only if Y is quasi-compact.
- (3) $\widetilde{s}_Y = \widetilde{s}_Z$ if and only if $\mathbf{Cl}^{\text{inv}}(Y) = \mathbf{Cl}^{\text{inv}}(Z)$.
- (4) $\widetilde{s}_Y = s_{\mathbf{Cl}^{\text{inv}}(Y)}$.

Remark 4.2. Note that the equivalence (1) of Proposition 4.1 can also be viewed in a topological way, since the sets of the type Y^{gen} , for Y varying among the subsets of $\text{Spec}(D)$, are the closed sets of a topology called the *Right-topology on $\text{Spec}(D)$* , which is the finest topology on $\text{Spec}(D)$ compatible with the opposite order of the given order on $\text{Spec}(D)$ [7, Lemma 2.1, Proposition 2.3(b)].

The following is a “finite type version” of Propositions 3.2, 3.4 and 3.5.

Proposition 4.3. *Let D be an integral domain, and denote by $\mathcal{I}_{\mathbf{f}}$ the set of nonzero finitely generated ideals of D . Let also $\widetilde{\Phi} : \mathbf{SStar}(D) \rightarrow \mathbf{SStar}_{\mathbf{f},sp}(D)$ be the map defined by $\widetilde{\Phi}(\star) := \widetilde{\star}$, for each $\star \in \mathbf{SStar}(D)$.*

- (1) *The set $\{\widetilde{U}_J := \mathbf{V}_J \cap \mathbf{SStar}_{\mathbf{f},sp}(D) \mid J \in \mathcal{I}_{\mathbf{f}}\}$ is a subbasis of open and quasi-compact subspaces for $\mathbf{SStar}_{\mathbf{f},sp}(D)$.*
- (2) *If $\mathbf{SStar}(D)$ is endowed with the Zariski topology, $\widetilde{\Phi}$ is a topological retraction.*
- (3) *If $\mathbf{SStar}(D)$ is endowed with the topology generated by the family $\{\mathbf{V}_J \mid J \in \mathcal{I}_{\mathbf{f}}\}$, then $\widetilde{\Phi}$ is the canonical map onto the Kolmogoroff quotient space of $\mathbf{SStar}(D)$.*
- (4) *If \star is a stable star operation of finite type on D , there is exactly one stable semistar operation of finite type \star on D such that $\star|_{\mathbf{F}(D)} = \star$.*
- (5) *The restriction $\lambda_{\mathbf{f}} : \mathbf{LS}_{\mathbf{f}}(D) \rightarrow \mathbf{SStar}_{\mathbf{f},sp}(D)$ of λ establishes a homeomorphism between spaces endowed with the Zariski topologies.*

Proof. The proofs follow essentially from the general case, with some additional care. For (1), we note that, for each $I \in \mathcal{I}$, $\mathbf{V}_I \cap \mathbf{SStar}_{\mathbf{f},sp}(D) = \bigcup \{\mathbf{V}_J \cap \mathbf{SStar}_{\mathbf{f},sp}(D) \mid J \subseteq I, J \in \mathbf{f}(D)\}$ (compare [13, Remark 2.2(d)]). To show that \widetilde{U}_J is quasi-compact when $J \in \mathcal{I}_{\mathbf{f}}$, consider the semistar operation s_Y , where $Y := \{P \in \text{Spec}(D) \mid P \not\subseteq J\}$. Since Y is quasi-compact, s_Y is of finite type (see [13, Corollary 4.4] or Proposition 4.1(2)). Moreover, $1 \in JD_P$ if and only if $P \in Y$; therefore, s_Y is the minimum of \widetilde{U}_J , and every open set containing s_Y contains the whole \widetilde{U}_J . It follows that \widetilde{U}_J is quasi-compact.

For (4), it is enough to note that if \star is of finite type then $\overline{\star}_e = \widetilde{\star}_e$, so that \star is of finite type if \star is. For (5), note that the image of $\lambda_{\mathbf{f}}$ is exactly $\mathbf{SStar}_{\mathbf{f},sp}(D)$ [14, Theorem 2.10, Corollary 2.11, and Proposition 3.2]. \square

The remaining part of the present section is devoted to the proof that $\mathbf{SStar}_{\mathbf{f},sp}(D)$ is a spectral space, in the sense of M. Hochster [24]. We start by studying the supremum and the infimum of a family of spectral operations.

Lemma 4.4. *Let \mathcal{D} be a nonempty set of spectral semistar operations. For each spectral semistar operation \star , set $\Delta(\star) := \text{QSpec}^*(D)$.*

- (1) $\wedge_{\mathcal{D}}$ is spectral with $\Delta(\wedge_{\mathcal{D}}) = \bigcup\{\Delta(\star) \mid \star \in \mathcal{D}\}$.
- (2) If $\vee_{\mathcal{D}}$ is quasi-spectral, then it is spectral with $\Delta(\vee_{\mathcal{D}}) = \bigcap\{\Delta(\star) \mid \star \in \mathcal{D}\}$.

Proof. (1) Set $\mathbf{\Delta} := \bigcup\{\Delta(\star) \mid \star \in \mathcal{D}\}$. For each $E \in \overline{\mathbf{F}}(D)$

$$E^{\wedge_{\mathcal{D}}} = \bigcap\{ED_P \mid P \in \mathbf{\Delta}, \star \in \mathcal{D}\} = \bigcap\{ED_P \mid P \in \mathbf{\Delta}\}.$$

In particular, $\wedge_{\mathcal{D}}$ is spectral and $\mathbf{\Delta} \subseteq \text{QSpec}^{\wedge_{\mathcal{D}}}(D)$. On the other hand, if $Q \in \text{QSpec}^{\wedge_{\mathcal{D}}}(D)$, then $Q^{\star} \neq D^{\star}$ for some $\star \in \mathcal{D}$, and this implies that $Q \in \text{QSpec}^*(D)$. Therefore, $\mathbf{\Delta} = \text{QSpec}^{\wedge_{\mathcal{D}}}(D)$.

(2) Let $P \in \text{QSpec}^{\vee_{\mathcal{D}}}(D)$. Then, P belongs to $\text{QSpec}^*(D)$ for each $\star \in \mathcal{D}$, i.e., $\text{QSpec}^{\vee_{\mathcal{D}}}(D) \subseteq \bigcap\{\Delta(\star) \mid \star \in \mathcal{D}\}$. Since each $\star \in \mathcal{D}$ is spectral, then, for each $E \in \overline{\mathbf{F}}(D)$, $E^{\star} \subseteq ED_P$ for all $P \in \Delta(\star)$, and in particular for all $P \in \bigcap\{\Delta(\star) \mid \star \in \mathcal{D}\}$. Hence,

$$E^{\vee_{\mathcal{D}}} \subseteq \bigcap\left\{ED_P \mid P \in \bigcap\{\Delta(\star) \mid \star \in \mathcal{D}\}\right\} \subseteq \bigcap\{ED_P \mid P \in \text{QSpec}^{\vee_{\mathcal{D}}}(D)\}.$$

However, if $\vee_{\mathcal{D}}$ is quasi-spectral, then it is known that the right hand side is contained in $E^{\vee_{\mathcal{D}}}$ [14, Proposition 4.8]. Therefore they are equal, and hence $\vee_{\mathcal{D}}$ is spectral, with $\Delta(\vee_{\mathcal{D}}) = \bigcap\{\Delta(\star) \mid \star \in \mathcal{D}\}$. \square

Example 4.5. In relation with Lemma 4.4(2), we note that the supremum of a family of spectral semistar operations may not be quasi-spectral. Indeed, let \mathbb{A} be the ring of algebraic integers, i.e., the integral closure of \mathbb{Z} in the algebraic closure $\overline{\mathbb{Q}}$ of \mathbb{Q} . Recall that \mathbb{A} is a one-dimensional Bézout domain [31, page 72].

Claim 1. *For each maximal ideal P of \mathbb{A} , $\text{Max}(\mathbb{A}) \setminus \{P\}$ is not a quasi-compact subspace of $\text{Max}(\mathbb{A})$ (endowed with the Zariski topology).*

By contradiction, since $\text{Max}(\mathbb{A}) \setminus \{P\}$ is open in $\text{Max}(\mathbb{A})$ it would be equal to $\text{D}(J) \cap \text{Max}(\mathbb{A})$ for some finitely generated ideal J of \mathbb{A} . Being \mathbb{A} a Bézout domain, this would imply that $\text{Max}(\mathbb{A}) \setminus \{P\} = \text{D}(\alpha) \cap \text{Max}(\mathbb{A})$ for some $\alpha \in \mathbb{A}$; in particular, the ideal $\alpha\mathbb{A}$ would be P -primary. Let K be the Galois closure of $\mathbb{Q}(\alpha)$ over \mathbb{Q} and consider the prime ideal $P_K := P \cap \mathcal{O}_K$, where \mathcal{O}_K the ring of integers of the field K . Let $P \cap \mathbb{Z} = p\mathbb{Z}$ for some prime integer p and let F be a Galois extension of \mathbb{Q} where p splits and such that $F \cap K = \mathbb{Q}$ (there are infinitely many such fields F , since p splits in infinitely many quadratic extensions of \mathbb{Q} and K contains only a finite number of them). We claim that P_K splits in the compositum FK : if this is true, then α would be contained in more than a single prime ideal of \mathbb{A} , against the hypothesis.

Set $P_{FK} := P \cap \mathcal{O}_{FK}$ and $P_F := P \cap \mathcal{O}_F = P_{FK} \cap \mathcal{O}_F$. Suppose P_K does not split in \mathcal{O}_{FK} : then $P_K \mathcal{O}_{FK}$ would be primary to P_{FK} . On the other hand, $P_F \cap \mathbb{Z} = p\mathbb{Z}$; since p splits in \mathcal{O}_F , and the Galois group of F over \mathbb{Q} acts transitively on the primes of \mathcal{O}_F lying over p , there is an automorphism σ of F such that $\sigma(P_F) \neq P_F$. Since $K \cap F = \mathbb{Q}$, there is an automorphism τ of FK such that $\tau|_F = \sigma$ and $\tau|_K$ is the identity. Therefore, $\tau(P_K) = P_K$ and $\tau(P_{FK})$ must contain P_K , i.e., $\tau(P_{FK}) = P_{FK}$. However, P_{FK} contains P_F , and $\tau(P_{FK})$ contains $\sigma(P_F)$; therefore, P_{FK} must contain both P_F and $\sigma(P_F)$, which is impossible. Therefore, P_K splits in \mathcal{O}_{FK} .

Claim 2. *For every $P \in \text{Max}(\mathbb{A})$, $\bigcap\{\mathbb{A}_Q \mid Q \in \text{Max}(\mathbb{A}) \setminus \{P\}\} = \mathbb{A}$.*

Let $B := \bigcap \{\mathbb{A}_Q \mid Q \in \text{Max}(\mathbb{A}) \setminus \{P\}\}$. By the previous claim, $\text{Max}(\mathbb{A}) \setminus \{P\}$ is not quasi-compact, and then it follows immediately that P belongs to the closure of $\text{Max}(\mathbb{A}) \setminus \{P\}$, with respect to the inverse topology. In other words, every maximal ideal of \mathbb{A} is a limit point in the inverse topology and so $\text{Max}(\mathbb{A})$ with the inverse topology is a perfect space. Finally, by [36, Proposition 5.6(4)], we have $B = \mathbb{A}$.

We are ready now to show that the supremum of a family of spectral semistar operations on \mathbb{A} may not be quasi-spectral. For every $P \in \text{Max}(\mathbb{A})$, let $\star_P := \mathfrak{s}_{\text{Max}(\mathbb{A}) \setminus \{P\}}$, and define $\star := \bigvee \{\star_P \mid P \in \text{Max}(\mathbb{A})\}$. By Claim 2, $\mathbb{A}^{\star_P} = \mathbb{A}$ for every $P \in \text{Max}(\mathbb{A})$, and thus $\mathbb{A}^\star = \mathbb{A}$. However, P is not a \star_P -ideal since $P^{\star_P} = \mathbb{A}$ and therefore $P^\star = \mathbb{A}$ for every $P \in \text{Max}(\mathbb{A})$. Since each nonzero principal (or, equivalently, finitely generated) integral ideal of \mathbb{A} is a \star -ideal and the set of nonzero prime \star -ideals of \mathbb{A} is empty, it follows that \star is not quasi-spectral.

Theorem 4.6. *Let D be an integral domain. The space $\text{SStar}_{f,sp}(D)$ of the stable semistar operations of finite type on D , endowed with the Zariski topology induced by $\text{SStar}(D)$, is a spectral space.*

Proof. In order to prove that a topological space X is a spectral space, we use the characterization given in [11, Corollary 3.3]. We recall that if \mathcal{B} is a nonempty family of subsets of X , for a given subset Y of X and an ultrafilter \mathcal{U} on Y , we set

$$Y_{\mathcal{B}}(\mathcal{U}) := \{x \in X \mid \text{for each } B \in \mathcal{B}, \text{ it happens that } x \in B \Leftrightarrow B \cap Y \in \mathcal{U}\}.$$

The subset Y of X is called \mathcal{B} -ultrafilter closed if $Y_{\mathcal{B}}(\mathcal{U}) \subseteq Y$; the \mathcal{B} -ultrafilter closed subsets of X are the closed subspaces of a topology on X called the \mathcal{B} -ultrafilter topology on X .

By [11, Corollary 3.3], for a topological space X being a spectral space is equivalent to X being a T_0 -space having a subbasis for the open sets \mathcal{S} such that $X_{\mathcal{S}}(\mathcal{U}) \neq \emptyset$, for each ultrafilter \mathcal{U} on X .

We already know that $\mathcal{X} := \text{SStar}_{f,sp}(D)$ is a T_0 -space. By Proposition 4.3(1), the collection of sets

$$\tilde{\mathcal{T}} := \{\tilde{\mathcal{U}}_J \mid J \subseteq D, J \text{ finitely generated ideal of } D\}$$

is a subbasis of the Zariski topology of \mathcal{X} . Let \mathcal{U} be any ultrafilter on \mathcal{X} ; the conclusion will follow if we prove that the set

$$\mathcal{X}_{\tilde{\mathcal{T}}}(\mathcal{U}) := \{\star \in \mathcal{X} \mid [\text{for each } \tilde{\mathcal{U}}_J \in \tilde{\mathcal{T}}, \text{ it happens that } \star \in \tilde{\mathcal{U}}_J \Leftrightarrow \tilde{\mathcal{U}}_J \in \mathcal{U}]\}$$

is nonempty.

Consider the semistar operation

$$\star := \bigvee \{\wedge_{\tilde{\mathcal{U}}_J} \mid \tilde{\mathcal{U}}_J \in \mathcal{U}\},$$

on D . By Lemma 4.4(1), each $\wedge_{\tilde{\mathcal{U}}_J}$ is spectral and, since $\tilde{\mathcal{U}}_J$ is quasi-compact (Proposition 4.3(1)), is also of finite type [13, Proposition 2.7]. By applying [3, p. 1628], it can be easily shown that \star is of finite type, and thus it is quasi-spectral [14, Corollary 4.21]. By Lemma 4.4(2), it follows that \star is spectral, i.e., $\star \in \mathcal{X}$.

To prove that $\star \in \mathcal{X}_{\tilde{\mathcal{T}}}(\mathcal{U})$, we apply the same argument used in proving [13, Theorem 2.13]: let $\tilde{\mathcal{U}}_J \in \mathcal{U}$. If $\star \in \tilde{\mathcal{U}}_J$, then by the definition of \star and [3, p.1628] (see also [13, Lemma 2.12]) there are finitely generated ideals F_1, F_2, \dots, F_n of D such that

$$1 \in J^{\wedge_{\tilde{\mathcal{U}}_{F_1}} \circ \wedge_{\tilde{\mathcal{U}}_{F_2}} \circ \dots \circ \wedge_{\tilde{\mathcal{U}}_{F_n}}},$$

with $\tilde{\mathfrak{U}}_{F_i} \in \mathcal{U}$. Each semistar operation $\sigma \in \tilde{\mathfrak{U}}_{F_1} \cap \tilde{\mathfrak{U}}_{F_2} \cap \cdots \cap \tilde{\mathfrak{U}}_{F_n}$ is bigger than each each semistar operation of type $\wedge_{\tilde{\mathfrak{U}}_{F_i}}$; it follows that $\tilde{\mathfrak{U}}_{F_1} \cap \tilde{\mathfrak{U}}_{F_2} \cap \cdots \cap \tilde{\mathfrak{U}}_{F_n} \subseteq \tilde{\mathfrak{U}}_J$, and thus the latter set is in \mathcal{U} . Conversely, if $\tilde{\mathfrak{U}}_J \in \mathcal{U}$, then $\star \geq \wedge_{\tilde{\mathfrak{U}}_J}$, and thus $1 \in J^\star$. The proof is now complete. \square

Remark 4.7. The proof of Theorem 4.6 actually shows more than just the fact that $\mathbf{SStar}_{f,sp}(D)$ is a spectral space. Given a spectral space X , the *constructible topology* on X is the coarsest topology such that every open and quasi-compact subset of X (in the original topology) is both open and closed. By [18], the closed sets of the constructible topology in X are the subsets Y of X such that, for every ultrafilter \mathcal{U} of Y ,

$$Y_{\mathcal{B}}(\mathcal{U}) := \{x \in X \mid \text{for each } B \in \mathcal{B}, \text{ it happens that } x \in B \Leftrightarrow B \cap Y \in \mathcal{U}\} \subseteq Y,$$

where \mathcal{B} is the set of open and quasi-compact subspaces of X . Therefore, in view of the proof of [13, Theorem 2.13], what we have actually proved in Theorem 4.6 is that, when $\mathbf{SStar}_f(D)$ is endowed with the constructible topology, $\mathbf{SStar}_{f,sp}(D)$ is a closed subspace.

From the previous theorem and Proposition 4.3(5), we deduce immediately the following:

Corollary 4.8. *Let D be an integral domain. The space $\mathbf{LS}_f(D)$ of the localizing systems of finite type on D , endowed with the Zariski topology induced by $\mathbf{LS}(D)$, is a spectral space.*

5. THE SPACE OF **eab** SEMISTAR OPERATIONS OF FINITE TYPE

A semistar operation \star on an integral domain D is said to be an ***eab** semistar operation* [respectively, an ***ab** semistar operation*] if, for every $F, G, H \in \mathbf{f}(D)$ [respectively, for every $F \in \mathbf{f}(D)$, $G, H \in \overline{\mathbf{F}}(D)$] the inclusion $(FG)^\star \subseteq (FH)^\star$ implies $G^\star \subseteq H^\star$. Note that, if \star is **eab**, then \star_f is also **eab**, since \star and \star_f agree on finitely generated fractional ideals. The concepts of **eab** and **ab** operations coincide on finite type operations, but not in general [19, 20].

Remark 5.1. W. Krull only considered the concept of “arithmetisch brauchbar” operation (for short **ab**-operation, as above) [32]. He did not consider the concept of “endlich arithmetisch brauchbar” operation (or, more simply, **eab**-operation as above), that instead stems from the original version of Gilmer’s book [22].

Let $\mathbf{Zar}(D) := \{V \mid V \text{ is a valuation overring of } D\}$ be equipped with the *Zariski topology*, i.e., the topology having, as subbasic open subspaces, the subsets $\mathbf{Zar}(D[x])$ for x varying in K . The set $\mathbf{Zar}(D)$, endowed with the Zariski topology, is often called the *Riemann-Zariski space of D* [42, Chapter VI, §17, page 110]. Recently, the use of Riemann-Zariski spaces had a strong impact on the study of algebraic properties of integrally closed domains. For a deeper insight on this topic see, for example, [37, 39, 36, 38].

A *valuative semistar operation* is a semistar operation of the type \wedge_Y , where $Y \subseteq \mathbf{Zar}(D)$; it is easy to see that it is an **eab** semistar operation. In particular, the **b**-operation, where $\mathfrak{b} := \wedge_{\mathbf{Zar}(D)}$, is an **eab** semistar operation of finite type on D , since $\mathbf{Zar}(D)$ is quasi-compact [13, Proposition 4.5]. More generally, for the same reason, for each overring T of D , the valuative semistar operation $\mathfrak{b}(T) := \wedge_{\mathbf{Zar}(T)}$ is an **eab** semistar operation of finite type on D .

Just like in the case of the relation between stable and spectral operations, not every **eab** semistar operation is valutative, but the two definitions agree on finite type operations (see, for instance, [15, Corollary 5.2]). However, unlike the spectral case, there are example of quasi-spectral **eab** operations that are not valutative [19, Example 15].

Denote by $\mathbf{SStar}_{\text{val}}(D)$ [respectively, $\mathbf{SStar}_{\text{eab}}(D)$; $\mathbf{SStar}_{f, \text{eab}}(D)$] the set of valutative [respectively, **eab**; **eab** of finite type] semistar operations on D , endowed with the Zariski topology induced from $\mathbf{SStar}(D)$. By the previous remarks, we have:

$$\mathbf{SStar}_{f, \text{eab}}(D) := \mathbf{SStar}_{\text{eab}}(D) \cap \mathbf{SStar}_f(D) = \mathbf{SStar}_{\text{val}}(D) \cap \mathbf{SStar}_f(D).$$

To every semistar operation $\star \in \mathbf{SStar}(D)$ we can associate a map $\star_a : \overline{\mathbf{F}}(D) \rightarrow \overline{\mathbf{F}}(D)$ defined by

$$F^{\star_a} := \bigcup \{((FG)^\star : G^\star) \mid G \in \mathbf{f}(D)\}$$

for every $F \in \mathbf{f}(D)$, and then extended to arbitrary modules $E \in \overline{\mathbf{F}}(D)$ by setting $E^{\star_a} := \bigcup \{F^{\star_a} \mid F \subseteq E, F \in \mathbf{f}(D)\}$. The map \star_a is always an **eab** semistar operation of finite type on D [15, Proposition 4.5(1, 2)]. Moreover, $\star = \star_a$ if and only if \star is an **eab** semistar operation of finite type, called *the eab semistar operation of finite type associated to \star* and, if \star is an **eab** semistar operation, then $\star_a = \star_f$ [15, Proposition 4.5(4)]. The following proposition is an analogue of Propositions 3.2(2) and 4.3(2).

Proposition 5.2. *Let D be an integral domain and let $\Phi_a : \mathbf{SStar}(D) \rightarrow \mathbf{SStar}_{f, \text{eab}}(D)$ be the map defined by $\Phi_a(\star) := \star_a$, for each $\star \in \mathbf{SStar}(D)$. Then, Φ_a is topological retraction of $\mathbf{SStar}(D)$ onto $\mathbf{SStar}_{f, \text{eab}}(D)$.*

Proof. We start by showing that Φ_a is continuous. Indeed, if H is a nonzero finitely generated fractional ideal of D ,

$$\begin{aligned} \Phi_a^{-1}(\mathbf{V}_H \cap \mathbf{SStar}_{f, \text{eab}}(D)) &= \{\star \in \mathbf{SStar}(D) \mid 1 \in H^{\star_a}\} \\ &= \{\star \in \mathbf{SStar}(D) \mid F^\star \subseteq (HF)^\star \text{ for some } F \in \mathbf{f}(D)\} \\ &= \bigcup \{\{\star \in \mathbf{SStar}(D) \mid F^\star \subseteq (HF)^\star\} \mid F \in \mathbf{f}(D)\}. \end{aligned}$$

Let $F := x_1D + x_2D + \cdots + x_nD$. Then, $F^\star \subseteq (HF)^\star$ is and only if $x_i \in (HF)^\star$ for $i = 1, \dots, n$, hence,

$$\Phi_a^{-1}(\mathbf{V}_H \cap \mathbf{SStar}_{f, \text{eab}}(D)) = \bigcup \left\{ \bigcap_{i=1}^n \mathbf{V}_{x_i^{-1}HF} \mid F \in \mathbf{f}(D) \right\},$$

which is an open set of $\mathbf{SStar}(D)$. Hence, Φ_a is continuous. Moreover, if \star is an **eab** operation of finite type, then $\star_a = (\star_a)_a$. Henceforth, Φ_a is a topological retraction. \square

We are now interested to what happens to the topological embedding ι , defined in Proposition 2.1, when restricted to subsets of integrally closed overrings. We start with a remark.

Remark 5.3. Let T be an overring of D , and let \star_T be a semistar operation on T . Then, we can define a semistar operation \star on D by $\star := \star_T \circ \wedge_{\{T\}}$, i.e., $E^\star := (ET)^{\star_T}$ for every $E \in \overline{\mathbf{F}}(D)$. If now $F \in \mathbf{f}(T)$, then

$$\begin{aligned} F^{\star_a} &= \bigcup \{((FG)^\star : G^\star) \mid G \in \mathbf{f}(D)\} = \bigcup \{((FGT)^{\star_T} : (GT)^{\star_T}) \mid G \in \mathbf{f}(D)\} = \\ &= \bigcup \{((FTH)^{\star_T} : H^{\star_T}) \mid H \in \mathbf{f}(T)\} = (FT)^{(\star_T)_a} = F^{(\star_T)_a}. \end{aligned}$$

Hence, for every $E \in \overline{\mathbf{F}}(D)$, $E^{\star a} = (ET)^{(\star_T)^a}$, that is, $\star_a = (\star_T)_a \circ \wedge_{\{T\}}$ (where $(\star_T)_a$ is the **eab** semistar operation of finite type on T associated to \star_T).

Olberding in [37, 38], considered a new topology (called the **b-topology**) on the set $\mathbf{Overr}_{ic}(D)$ of integrally closed overrings of D by taking, as a subbasis of open sets, the sets of the form

$$\mathcal{U}_{ic}(F, G) := \{T \in \mathbf{Overr}_{ic}(D) \mid F \subseteq G^{b(T)}\},$$

where F and G range among the nonzero finitely generated D -submodules of K . He showed that the **b-topology** on $\mathbf{Overr}_{ic}(D)$ is finer than (or equal to) the Zariski topology (since it is straightforward that $\mathbf{B}_F = \mathcal{U}_{ic}(F, D)$, where $\mathbf{B}_F := \mathbf{Overr}_{ic}(D[F]) = \mathbf{Overr}_{ic}(D[x_1, x_2, \dots, x_n])$, for each $F = x_1D + x_2D + \dots + x_nD \in \mathbf{f}(D)$, is subbasic open set of $\mathbf{Overr}_{ic}(D)$ with the topology induced by the Zariski topology of $\mathbf{Overr}(D)$) and the two topologies coincide when restricted to the Riemann-Zariski space $\text{Zar}(D)$ [37, Corollary 2.8]. Using semistar operations, we can show more, i.e, the **b-topology** and the Zariski topology coincide on $\mathbf{Overr}_{ic}(D)$ (Corollary 5.5).

Proposition 5.4. *Let D be an integral domain, and consider the injective map $\iota_{ic,a} : \mathbf{Overr}_{ic}(D) \rightarrow \mathbf{SStar}_f(D)$ defined by $\iota_{ic,a}(T) := \mathbf{b}(T)$, for each $T \in \mathbf{Overr}_{ic}(D)$. Assume that $\mathbf{SStar}_f(D)$ is endowed with the Zariski topology. Then:*

- (1) *If $\mathbf{Overr}_{ic}(D)$ is endowed with the Zariski topology, then $\iota_{ic,a}$ is continuous and injective.*
- (2) *If $\mathbf{Overr}_{ic}(D)$ is endowed with the **b-topology**, then $\iota_{ic,a}$ is topological embedding, i.e., it establishes a homeomorphism between $\mathbf{Overr}_{ic}(D)$ and its image.*

Proof. (1) Let $\Psi_a := \Phi_a|_{\mathbf{SStar}_f(D)} : \mathbf{SStar}_f(D) \rightarrow \mathbf{SStar}_{f, \text{eab}}(D)$, defined by $\Psi_a(\star) = \star_a$, for each $\star \in \mathbf{SStar}_f(D)$ (Lemma 5.2). Note that $\iota_{ic,a} = \Psi_a \circ \iota$, since by Remark 5.3 we have $(\wedge_{\{T\}})_a = \mathbf{b}(T)$ for each overring T of D (see also the proof of Proposition 5.6). Therefore, $\iota_{ic,a}$ is continuous as a composition of continuous maps. Moreover, if $T', T'' \in \mathbf{Overr}_{ic}(D)$ and $T' \neq T''$, then $\text{Zar}(T') \neq \text{Zar}(T'')$ and so $\mathbf{b}(T') \neq \mathbf{b}(T'')$, i.e., $\iota_{ic,a}$ is injective.

(2) Since the **b-topology** is finer than the Zariski topology, $\iota_{ic,a}$ is continuous by the previous part of the proof. We set

$$\mathbf{SStar}_b(D) := \{\mathbf{b}(T) \mid T \in \mathbf{Overr}_{ic}(D)\} = \iota_{ic,a}(\mathbf{Overr}_{ic}(D)).$$

Let $\mathcal{U}_{ic}(F, G)$ be a subbasic open set of $\mathbf{Overr}_{ic}(D)$ in the **b-topology**. If $F := x_1D + x_2D + \dots + x_nD$, then $\mathcal{U}_{ic}(F, G) = \mathcal{U}_{ic}(x_1, G) \cap \mathcal{U}_{ic}(x_2, G) \cap \dots \cap \mathcal{U}_{ic}(x_n, G)$, so we can suppose that $F = xD$ for some $0 \neq x \in K$. In this situation we have:

$$\iota_{ic,a}(\mathcal{U}_{ic}(xD, G)) = \{\star \in \mathbf{SStar}_b(D) \mid x \in G^\star\} = \mathbf{V}_{x^{-1}G} \cap \mathbf{SStar}_b(D).$$

Therefore, $\iota_{ic,a}$ is open onto $\mathbf{SStar}_b(D)$ and hence it establishes a homeomorphism between $\mathbf{Overr}_{ic}(D)$ (with the **b-topology**) and $\mathbf{SStar}_b(D)$ (with the Zariski topology). \square

Corollary 5.5. *The **b-topology** and the Zariski topology coincide on $\mathbf{Overr}_{ic}(D)$.*

Proof. With the notation of the proof of Proposition 5.4(2), it is enough to observe that $\mathcal{U}_{ic}(xD, G)$ is also open in the Zariski topology of $\mathbf{Overr}_{ic}(D)$, since $\iota_{ic,a}^{-1}(\mathbf{V}_{x^{-1}G}) = \mathcal{U}_{ic}(xD, G)$. \square

The following can be seen as a companion of Proposition 3.1.

Proposition 5.6. *Let D be an integral domain and let $T \in \mathbf{Overr}(D)$. The following properties are equivalent:*

- (i) $\wedge_{\{T\}} \in \mathbf{SStar}_{f, \text{eab}}(D)$;
- (ii) $\iota(T) = \iota_{ic, a}(T)$;
- (iii) T is a Prüfer domain.

Proof. By definition of the **eab** semistar operation of finite type associated to $\wedge_{\{T\}}$, we have that $\wedge_{\{T\}} \in \mathbf{SStar}_{f, \text{eab}}(D)$ if and only if $\wedge_{\{T\}} = (\wedge_{\{T\}})_a$. However, by Remark 5.3 and Proposition 5.4, and noting that $\wedge_{\{T\}}$ (when restricted to $\overline{\mathbf{F}}(T)$) coincides with the identity semistar \mathbf{d}_T on $\overline{\mathbf{F}}(T)$, we have

$$(\wedge_{\{T\}})_a = (\mathbf{d}_T)_a \circ \wedge_{\{T\}} = \mathbf{b}_T \circ \wedge_{\{T\}} = \mathbf{b}(T) = \iota_{ic, a}(T),$$

where \mathbf{b}_T is the **b**-operation on T and $\mathbf{b}(T) = \wedge_{\text{Zar}(T)}$. Therefore, (i) and (ii) are equivalent. It is obvious that (iii) \Rightarrow (ii). For the reverse implication, it is enough to note that $\mathbf{d}_T = \mathbf{b}_T$ is equivalent to T being Prüfer [21, Lemma 2]. \square

Note that we can define the **b**-topology also on the whole $\mathbf{Overr}(D)$. The properties that we obtain are somewhat similar to Propositions 3.2(3) and 4.3(3).

Proposition 5.7. *Let D be an integral domain, and define the map $\beta : \mathbf{Overr}(D) \rightarrow \mathbf{Overr}_{ic}(D)$ by setting $\beta(T) := T^b = \overline{T}$, where \overline{T} is the integral closure of T .*

- (1) *If $\mathbf{Overr}_{ic}(D)$ and $\mathbf{Overr}(D)$ are endowed with the Zariski topology, then β is continuous and, hence, it is a topological retraction of $\mathbf{Overr}(D)$ onto $\mathbf{Overr}_{ic}(D)$.*
- (2) *If $\mathbf{Overr}_{ic}(D)$ is endowed with the Zariski topology and $\mathbf{Overr}(D)$ is endowed with the **b**-topology, then β is the canonical map onto the Kolmogoroff quotient space of $\mathbf{Overr}(D)$.*

Proof. (1) Let x be a nonzero element of K and let \mathbf{B}_x be a subbasic open set of the Zariski topology on $\mathbf{Overr}_{ic}(T)$. Then,

$$\begin{aligned} \beta^{-1}(\mathbf{B}_x) &= \{T \in \mathbf{Overr}(D) \mid x \in \overline{T}\} = \{T \in \mathbf{Overr}(D) \mid x \text{ is integral over } T\} = \\ &= \bigcup \{ \mathbf{B}_{\{\alpha_{k-1}, \alpha_{k-2}, \dots, \alpha_0\}} \mid x^k + \alpha_{k-1}x^{k-1} + \dots + \alpha_1x + \alpha_0 = 0 \} \end{aligned}$$

which is open since it is a union of open sets. Hence, β is continuous. It is a retraction since, for $T \in \mathbf{Overr}_{ic}(D)$, $T^b = T$.

(2) Since the Zariski topology on $\mathbf{Overr}_{ic}(D)$ coincides with the **b**-topology, we can consider on $\mathbf{Overr}_{ic}(D)$ the subbasic open sets $\mathcal{U}_{ic}(F, G)$, for $F, G \in \mathbf{f}(D)$. However, since $b(T) = b(\overline{T})$, we have $\beta^{-1}(\mathcal{U}_{ic}(F, G)) = \mathcal{U}(F, G) := \{T \in \mathbf{Overr}(D) \mid F \subseteq G^{b(T)}\}$, and thus β is continuous. In the same way, $\beta(\mathcal{U}(F, G)) = \mathcal{U}_{ic}(F, G)$, so β is open. Finally, it is easy to see that two overrings T' and T'' are topologically distinguishable by the **b**-topology if and only if they have the same integral closure; hence, β is the canonical map onto the Kolmogoroff quotient space of $\mathbf{Overr}(D)$ with the **b**-topology. \square

The relation between valutative operations and subsets of $\text{Zar}(D)$ exhibits a similar behaviour to the relation between spectral operations and subsets of $\text{Spec}(D)$. Similarly to the prime spectrum case (see the paragraph preceding Proposition 4.1), for each subset Y of the Riemann-Zariski space $\text{Zar}(D)$, we set $Y^{\text{gen}} := \{z \in \text{Zar}(D) \mid y \in \mathbf{Cl}(\{z\}) \text{ for some } y \in Y\}$ and we denote $\mathbf{Cl}^{\text{inv}}(Y)$ the closure of Y in

the inverse topology of the spectral space $\text{Zar}(D)$ (see [6], [24, Proposition 8] and [12, 13]). Compare the next lemma with Proposition 4.1.

Lemma 5.8. *Let D be an integral domain and let Y and Z be two nonempty subsets of $\text{Zar}(D)$. Then, the following statements hold.*

- (1) $\wedge_Y = \wedge_Z$ if and only if $Y^{\text{gen}} = Z^{\text{gen}}$.
- (2) \wedge_Y is of finite type if and only if Y is quasi-compact.
- (3) $(\wedge_Y)_f = (\wedge_Z)_f$ if and only if $\mathbf{Cl}^{\text{inv}}(Y) = \mathbf{Cl}^{\text{inv}}(Z)$.
- (4) $(\wedge_Y)_f = \wedge_{\mathbf{Cl}^{\text{inv}}(Y)}$.

Proof. (1) Note that, in the present situation, $Y^{\text{gen}} = \{V \in \text{Zar}(D) \mid V \supseteq V_0, \text{ for some } V_0 \in Y\}$. Assume first that $\wedge_Y = \wedge_Z$. Let V be a valuation domain such that $V \in Y^{\text{gen}} \setminus Z^{\text{gen}}$. Then, for any $W \in Z$, we can pick an element $x_W \in W \setminus V$. It follows that $I := (x_W^{-1} \mid W \in Z) \subseteq M_V$, where M_V is the maximal ideal of V . Thus, if $V_0 \in Y$ is such that $V_0 \subseteq V$ (such a V_0 exists since $V \in Y^{\text{gen}}$), we have $IV_0 \subseteq M_{V_0}$ and, in particular, $1 \notin I^{\wedge_Y}$. On the other hand, clearly $1 \in I^{\wedge_Z}$, a contradiction. The converse it is straightforward since, for each $Y \subseteq \text{Zar}(D)$, $\wedge_Y = \wedge_{Y^{\text{gen}}}$.

For (2), (3) and (4), see respectively [13, Proposition 4.5], [12, Theorem 4.9] and [12, Corollary 4.17]. \square

Remark 5.9. (a) Since $\mathfrak{b} = \wedge_{\text{Zar}(D)}$ is a semistar operation of finite type on D (and this can be proved completely independently from the topological point of view, see [30, Proposition 6.8.2] and [13, Remark 4.6]), from Lemma 5.8 we get a new proof of the fact that $\text{Zar}(D)$ is a quasi-compact space (this is a special case of Zariski's theorem [42, Theorem 40, page 113]).

(b) Note that the equivalence (1) of Lemma 5.8 can also be viewed in a topological way, as indicated in Remark 4.2, after replacing $\text{Zar}(D)$ to $\text{Spec}(D)$.

Using the \mathfrak{b} -operation, Krull introduced a general version of the classical Kronecker function ring, coinciding in case of Dedekind domains with the classical one (considered by L. Kronecker [41, 8, 23]). In fact, a Kronecker function ring can be defined starting by any \mathfrak{eab} semistar operation. In the next lemma, we summarize some properties of the Kronecker function ring, relevant to the remaining part of the paper.

Lemma 5.10. *Let D be an integral domain, \star an \mathfrak{eab} semistar operation on D , \mathcal{X} an indeterminate over D and let $\mathfrak{c}(h)$ be the content of a polynomial $h \in D[\mathcal{X}]$ (i.e., the ideal of D generated by the coefficients of h). Set $\mathcal{V}(\star) := \{V \in \text{Zar}(D) \mid F^\star \subseteq FV, \text{ for each } F \in \mathfrak{f}(D)\}$ and*

$$\text{Kr}(D, \star) := \{f/g \mid f, g \in D[\mathcal{X}], g \neq 0, \text{ with } \mathfrak{c}(f) \subseteq \mathfrak{c}(g)^\star\}.$$

Denote by $V(\mathcal{X})$ the Gaussian (or trivial) extension of V to $K(\mathcal{X})$ (i.e., $V(\mathcal{X}) = V[\mathcal{X}]_{M[\mathcal{X}]}$, where M is the maximal ideal of V).

- (1) $\text{Kr}(D, \star)$ is a Bézout domain with quotient field $K(\mathcal{X})$, called the \star -Kronecker function ring of D and, for each polynomial $f \in D[\mathcal{X}]$, $\mathfrak{c}(f)\text{Kr}(D, \star) = f\text{Kr}(D, \star)$.
- (2) $\text{Kr}(D, \star) = \bigcap \{V(\mathcal{X}) \mid V \in \mathcal{V}(\star)\}$ and, for each $E \in \overline{\mathbf{F}}(D)$, $E^{\star a} = E\text{Kr}(D, \star) \cap K = \bigcap \{EV \mid V \in \mathcal{V}(\star)\}$.
- (3) If T is an overring of D , then $\mathcal{V}((\wedge_{\{T\}})_a) = \text{Zar}(T)$; in particular, we reobtain that $(\wedge_{\{T\}})_a = \mathfrak{b}(T)$.

Proof. For (1), see [15, Definition 3.2, Propositions 3.3 and 3.11(2), Corollary 3.4(2) and Theorem 5.1], [23, Theorem 32.11] and [17, Theorems 11 and 14]. For the proof of (2) see [15, Proposition 4.1(5)] and [17, Theorem 14]. (3) is a direct consequence of (2) and of the definitions. \square

When considering Kronecker function rings, particularly important is the case $\star = \mathfrak{b}$ and, in this case, we simply set $\text{Kr}(D) := \text{Kr}(D, \mathfrak{b})$. In this situation, it follows easily from the fact that $\text{Kr}(D)$ is a Bézout domain that the localization map $\text{Spec}(\text{Kr}(D)) \rightarrow \text{Zar}(\text{Kr}(D))$ (defined by $P \mapsto \text{Kr}(D)_P$) is actually a homeomorphism. Moreover, the map $\Psi : \text{Zar}(D) \rightarrow \text{Zar}(\text{Kr}(D))$ (defined by $V \mapsto V(\mathbb{X})$) is a homeomorphism too [12, Propositions 3.1 and 3.3], so that, by appropriate compositions, we deduce that there is a canonical homeomorphism between $\text{Spec}(\text{Kr}(D))$ and $\text{Zar}(D)$.

We are now in condition to prove the main result of this section.

Theorem 5.11. *Let D be an integral domain with quotient field K . Let $\text{Kr}(D) := \text{Kr}(D, \mathfrak{b})$ be the \mathfrak{b} -Kronecker function ring of D and let $\theta : \text{Spec}(\text{Kr}(D)) \rightarrow \text{Zar}(D)$ be the homeomorphism defined by $\theta(Q) := \text{Kr}(D)_Q \cap K$, for each $Q \in \text{Spec}(\text{Kr}(D))$ [6, Theorem 2]. Then,*

- (1) *The homeomorphism θ induces a continuous bijection $\Theta : \text{SStar}_{sp}(\text{Kr}(D)) \rightarrow \text{SStar}_{val}(D)$ defined by setting $\Theta(s_{\mathcal{Y}}) := \wedge_{Z(\mathcal{Y})}$, for each $\mathcal{Y} \subseteq \text{Spec}(\text{Kr}(D))$, where $Z(\mathcal{Y}) := \{V \in \text{Zar}(D) \mid M_V(\mathbb{X}) \cap \text{Kr}(D) \in \mathcal{Y}\}$ and M_V is the maximal ideal of V .*
- (2) *The map Θ , restricted to the semistar operations of finite type, gives rise to a homeomorphism $\Theta_f : \text{SStar}_{f,sp}(\text{Kr}(D)) \rightarrow \text{SStar}_{f, eab}(D)$ of topological spaces (endowed with the Zariski topology).*
- (3) *$\text{SStar}_{f, eab}(D)$ is a spectral space.*

Proof. (1) It is straightforward that the homeomorphism θ from $\text{Spec}(\text{Kr}(D))$ to $\text{Zar}(D)$ (which is, in particular, a isomorphism of partially ordered sets with the ordering induced by their topologies) induces a 1-1 correspondence Θ_0 between the set $\{\mathcal{Y} \subseteq \text{Spec}(\text{Kr}(D)) \mid \mathcal{Y} = \mathcal{Y}^\downarrow\}$ (where $\mathcal{Y}^\downarrow := \{Q \in \text{Spec}(\text{Kr}(D)) \mid Q \subseteq Q', \text{ for some } Q' \in \mathcal{Y}\} = \mathcal{Y}^{\text{gen}}\}$ and the set $\{Y \subseteq \text{Zar}(D) \mid Y = Y^\uparrow\}$ (where $Y^\uparrow := \{W \in \text{Zar}(D) \mid W \supseteq W', \text{ for some } W' \in Y\} = Y^{\text{gen}}\}$. Therefore Θ_0 induces a bijection $\Theta : \text{SStar}_{sp}(\text{Kr}(D)) \rightarrow \text{SStar}_{val}(D)$ defined by $\Theta(s_{\mathcal{Y}}) := \wedge_{\Theta_0(\mathcal{Y})}$, where $\Theta_0(\mathcal{Y}) = \{V \in \text{Zar}(D) \mid M_V(\mathbb{X}) \cap \text{Kr}(D) \in \mathcal{Y}\} =: Z(\mathcal{Y})$ (cf. Lemma 5.8, [13, Corollaries 4.4 and 5.2, Proposition 5.1] and [14, Lemma 4.2 and Remark 4.5]).

Moreover, the map Θ is continuous by [13, Proposition 3.1(1)], since $(E\text{Kr}(D))^{s_{\mathcal{Y}}} \cap K = E^{\wedge_{Z(\mathcal{Y})}}$ for each $E \in \overline{\mathbf{F}}(D)$ and for each $\mathcal{Y} \subseteq \text{Spec}(\text{Kr}(D))$. In fact,

$$\begin{aligned} E^{\wedge_{Z(\mathcal{Y})}} &= \bigcap \{EV \mid V \in Z(\mathcal{Y})\} = \bigcap \{EV(\mathbb{X}) \cap K \mid V \in Z(\mathcal{Y})\} \\ &= E(\bigcap \{V(\mathbb{X}) \mid V \in Z(\mathcal{Y})\}) \cap K = E(\text{Kr}(D))^{s_{\mathcal{Y}}} \cap K \\ &= (E\text{Kr}(D))^{s_{\mathcal{Y}}} \cap K. \end{aligned}$$

(2) Since quasi-compact sets correspond biunivocally to finite type semistar operations, in both spectral and the valutive case, then Θ restricts to a continuous bijection $\Theta_f : \text{SStar}_{f,sp}(\text{Kr}(D)) \rightarrow \text{SStar}_{f, eab}(D)$ (see also [13, Proposition 3.1(2)]).

Let J be a nonzero finitely generated ideal of $\text{Kr}(D)$, thus it is principal (since $\text{Kr}(D)$ is a Bézout domain). Therefore $J = z\text{Kr}(D)$, for some nonzero element $z := \alpha/\beta \in K(\mathbb{X})$, where $\alpha, \beta \in D[\mathbb{X}]$ and α, β are nonzero. We can consider the basic open set $V_{\alpha/\beta}^\# := \{\star \in \text{SStar}(\text{Kr}(D)) \mid 1 \in ((\alpha/\beta)\text{Kr}(D))^\star\}$ of $\text{SStar}(\text{Kr}(D))$ (the

superscript \sharp is used here to emphasize the fact that we are considering subspaces of $\mathbf{SStar}(\mathbf{Kr}(D))$ and not of $\mathbf{SStar}(D)$.

$$\text{Claim. } \Theta_f \left(\mathbf{V}_{\alpha/\beta}^\sharp \cap \mathbf{SStar}_{f,sp}(\mathbf{Kr}(D)) \right) = \left(\bigcap_{i=1}^n \mathbf{V}_{b_i^{-1}\mathbf{c}(\alpha)} \right) \cap \mathbf{SStar}_{f,eab}(D),$$

(where $\beta := b_0 + b_1\mathbb{X} + \dots + b_n\mathbb{X}^n$ and $\mathbf{c}(\alpha)$ is the ideal of D generated by the coefficients of the polynomial α).

Indeed, let $\tilde{\mathbf{U}}_{\alpha/\beta}^\sharp := \mathbf{V}_{\alpha/\beta}^\sharp \cap \mathbf{SStar}_{f,sp}(\mathbf{Kr}(D))$, suppose that $\star \in \tilde{\mathbf{U}}_{\alpha/\beta}^\sharp$ and let $\ast := \Theta_f(\star)$. Then, $1 \in (\alpha/\beta \cdot \mathbf{Kr}(D))^\star$, i.e., $(\beta\mathbf{Kr}(D))^\star \subseteq (\alpha\mathbf{Kr}(D))^\star$. However, $\beta\mathbf{Kr}(D) = \mathbf{c}(\beta)\mathbf{Kr}(D)$ (Lemma 5.10(1)), and analogously for α ; thus,

$$\mathbf{c}(\beta) \subseteq (\mathbf{c}(\beta)\mathbf{Kr}(D))^\star \cap K \subseteq (\mathbf{c}(\alpha)\mathbf{Kr}(D))^\star \cap K = \mathbf{c}(\alpha)^\star.$$

Hence, $b_i \in \mathbf{c}(\alpha)^\star$ for each i , and $1 \in (b_i^{-1}\mathbf{c}(\alpha))^\star$, that is, $\ast \in \mathbf{V}_{b_i^{-1}\mathbf{c}(\alpha)}$ for every i . On the other hand, it is a straightforward consequence of the definition that $\ast \in \mathbf{SStar}_{f,eab}(D)$.

Conversely, let $\ast \in (\bigcap_{i=1}^n \mathbf{V}_{b_i^{-1}\mathbf{c}(\alpha)}) \cap \mathbf{SStar}_{f,eab}(D)$. Since Θ_f is bijective, then $\ast = \Theta_f(\star)$ for a unique $\star \in \mathbf{SStar}_{f,sp}(\mathbf{Kr}(D))$. Then, $b_i \in \mathbf{c}(\alpha)^\star$ for every i , and $\mathbf{c}(\beta) \subseteq \mathbf{c}(\alpha)^\star$; it follows that $\beta/\alpha \in \mathbf{Kr}(D, \ast)$, i.e., $1 \in \alpha/\beta \cdot \mathbf{Kr}(D, \ast)$. On the other hand, $\ast = \Theta_f(\star)$ implies $E^\ast = (E\mathbf{Kr}(D))^\star \cap K$ and, moreover, since \ast is **eab**, $E^\ast = E\mathbf{Kr}(D, \ast) \cap K$, for each $E \in \overline{\mathbf{F}}(D)$ (Lemma 5.10(2)). Therefore, $1 \in \alpha/\beta \cdot \mathbf{Kr}(D, \ast)$ implies that $1 \in (\alpha/\beta \cdot \mathbf{Kr}(D))^\star$, i.e., $\star \in \tilde{\mathbf{U}}_{\alpha/\beta}^\sharp$, so that $\ast \in \Theta_f(\tilde{\mathbf{U}}_{\alpha/\beta}^\sharp)$.

The claim ensures that Θ_f is open, and hence we conclude that it is a homeomorphism.

(3) is an easy consequence of (2) and Theorem 4.6. \square

Remark 5.12. As we did in Remark 4.7, we can ask if $\mathbf{SStar}_{f,eab}(D)$ is closed when $\mathbf{SStar}_f(D)$ is endowed with the constructible topology. The answer is positive; indeed, consider the map $\mathbf{\Lambda} : \mathbf{SStar}_{f,sp}(\mathbf{Kr}(D)) \rightarrow \mathbf{SStar}_f(D)$ obtained by composing Θ_f with the inclusion of $\mathbf{SStar}_{f,eab}(D)$ into $\mathbf{SStar}_f(D)$. Let $I \in \mathbf{f}(D)$ and let \mathbf{V}_I be a subbasic open set of $\mathbf{SStar}(D)$ then, by the proof of Theorem 5.11, there exist $\alpha, \beta \in D[\mathbb{X}]$, with α, β nonzero, such that

$$\mathbf{\Lambda}^{-1}(\mathbf{V}_I \cap \mathbf{SStar}_f(D)) = \mathbf{V}_{\alpha/\beta}^\sharp \cap \mathbf{SStar}_{f,sp}(\mathbf{Kr}(D)) = \tilde{\mathbf{U}}_{((\alpha/\beta)\mathbf{Kr}(D)) \cap \mathbf{Kr}(D)}^\sharp.$$

Since $((\alpha/\beta)\mathbf{Kr}(D)) \cap \mathbf{Kr}(D)$ is finitely generated ($\mathbf{Kr}(D)$ being a Prüfer domain [23, (25.4), part (1)]), $\mathbf{\Lambda}^{-1}(\mathbf{V}_I \cap \mathbf{SStar}_f(D))$ is quasi-compact (Proposition 4.3(1)). This means that $\mathbf{\Lambda}$ is a spectral map. In particular, it is continuous when $\mathbf{SStar}_{f,sp}(\mathbf{Kr}(D))$ and $\mathbf{SStar}_f(D)$ are endowed with the constructible topology. Since a spectral space, endowed with the constructible topology, is both compact and Hausdorff, $\mathbf{\Lambda}$ is a closed map, when both spaces are endowed with the constructible topology. In particular, $\mathbf{\Lambda}(\mathbf{SStar}_{f,sp}(\mathbf{Kr}(D))) = \mathbf{SStar}_{f,eab}(D)$ is closed subspace of $\mathbf{SStar}_f(D)$, endowed with the constructible topology.

Acknowledgment. We sincerely thank the referee for the careful reading of the manuscript and for providing constructive comments and help in improving the presentation of the paper.

REFERENCES

- [1] D.D. Anderson, Star-operations induced by overrings, *Comm. Algebra* **16** (1988), 2535–2553.
- [2] D.D. Anderson and S.J. Cook, Two star operations and their induced lattices, *Comm. Algebra* **28** (2000), 2461–2475.
- [3] D.F. Anderson and D.D. Anderson, Examples of star operations on integral domains, *Comm. Algebra* **18** (1990), 1621–1643.
- [4] N. Bourbaki, *Algèbre Commutative*, Chap. 1-2, Hermann, Paris, 1961.
- [5] S.T. Chapman and S. Glaz, One hundred problems in commutative ring theory, in “Non-Noetherian Commutative Ring Theory”, Kluwer Academic Publishers, Dordrecht, 2000.
- [6] D. Dobbs and M. Fontana, Kronecker function rings and abstract Riemann surfaces. *J. Algebra* **99** (1986), 263–274.
- [7] D. Dobbs, M. Fontana, I. Papick, On certain distinguished spectral sets. *Ann. Mat. Pura Appl.* (4) **128** (1981), 227–240.
- [8] H. M. Edwards, *Divisor Theory*. Birkhäuser, Boston, 1990.
- [9] N. Epstein, A guide to closure operations in commutative algebra. *Progress in Commutative Algebra* 2, 1–37, Walter de Gruyter, Berlin, 2012.
- [10] N. Epstein, Semistar operations and standard closure operations, *Comm. Algebra* **43** (2015), 325–336.
- [11] C. A. Finocchiaro, Spectral spaces and ultrafilters, *Comm. Algebra* **42** (2014), 1496–1508.
- [12] C. A. Finocchiaro, M. Fontana, and K. A. Loper, The constructible topology on spaces of valuation domains, *Trans. Am. Math. Soc.* (2013), 6199–6216.
- [13] C. A. Finocchiaro and D. Spirito, Some topological considerations on semistar operations, *J. Algebra*, **409**(2014), 199–218.
- [14] M. Fontana and J. Huckaba, Localizing systems and semistar operations, in “Non-Noetherian Commutative Ring Theory” (Scott T. Chapman and Sarah Glaz, eds.), Kluwer Academic Publishers, 2000, pp. 169–198.
- [15] M. Fontana and K.A. Loper, Kronecker function rings: a general approach, in “Ideal theoretic methods in commutative algebra” (Columbia, MO, 1999), 189–205, Lecture Notes in Pure and Appl. Math., **220**, Dekker, New York, 2001.
- [16] M. Fontana and K.A. Loper, Nagata rings, Kronecker function rings and related semistar operations, *Comm. Algebra* **31** (2003), 4775–4805
- [17] M. Fontana and K.A. Loper, An historical overview of Kronecker function rings, Nagata rings, and related star and semistar operations, in “*Multiplicative Ideal Theory in Commutative Algebra: A Tribute to the Work of Robert Gilmer*” (J.W. Brewer, S. Glaz, W. Heinzer, and B. Olberding Editors), 169–187, Springer, New York 2006.
- [18] M. Fontana and K.A. Loper, The patch topology and the ultrafilter topology on the prime spectrum of a commutative ring, *Comm. Algebra* **36** (2008), 2917–2922.
- [19] M. Fontana and K. A. Loper, Cancellation properties in ideal systems: a classification of e.a.b. semistar operations, *J. Pure Appl. Algebra* **213** (2009), 2095–2103.
- [20] M. Fontana, K. A. Loper, and R. Matsuda, Cancellation properties in ideal systems: an e.a.b. not a.b. star operation, *AJSE (Arabian Journal for Science and Engineering)–Mathematics*, **35** (2010), 45–49.
- [21] M. Fontana and G. Picozza, On some classes of integral domains defined by Krull’s a.b. operations. *J. Algebra* **341** (2011), 179–197.
- [22] R. Gilmer, *Multiplicative Ideal Theory*, Vol. I & II, Queen’s Papers in Pure and Applied Mathematics, Kingston, Ontario, Canada, 1968.
- [23] R. Gilmer, *Multiplicative Ideal Theory*, M. Dekker, New York, 1972.
- [24] M. Hochster, Prime ideal structure in commutative rings, *Trans. Amer. Math. Soc.* **142** (1969), 43–60.
- [25] J. Huckaba, *Commutative rings with zero divisors*, M. Dekker, New York, 1988.
- [26] F. Halter-Koch, *Ideal systems. An introduction to multiplicative ideal theory*. Monographs and Textbooks in Pure and Applied Mathematics, **211** Marcel Dekker, Inc., New York, 1998.
- [27] F. Halter-Koch, Construction of ideal systems with nice Noetherian properties. *Non-Noetherian commutative ring theory*, 271–285, Math. Appl., **520**, Kluwer Acad. Publ., Dordrecht, 2000.
- [28] F. Halter-Koch, Localizing systems, module systems, and semistar operations, *J. Algebra* **238** (2001), 723–761.

- [29] F. Halter-Koch, Multiplicative ideal theory in the context of commutative monoids. *Commutative algebra: Noetherian and non-Noetherian perspectives*, 203–231, Springer, New York, 2011.
- [30] C. Huneke and I. Swanson, Integral Closure of Ideals, Rings, and Modules, London Math. Soc. Lecture Note Ser., vol. 336, Cambridge University Press, Cambridge, 2006.
- [31] I. Kaplansky, Commutative rings, Allyn and Bacon Boston, 1970.
- [32] W. Krull, Beiträge zur Arithmetik kommutativer Integritätsbereiche, I - II. *Math. Z.* **41** (1936), 545–577; 665–679.
- [33] W. Krull, Gesammelte Abhandlungen / Collected Papers, Hrsg. v. Paulo Ribenboim, Walter de Gruyter, Berlin, 1999.
- [34] H. Matsumura, Commutative ring theory, Cambridge University Press, Cambridge, 1986.
- [35] A. Okabe and R. Matsuda, Semistar operations on integral domains, *Math. J. Toyama Univ.* **17** (1994), 1–21.
- [36] B. Olberding, Affine schemes and topological closures in the Zariski-Riemann space of valuation rings, *J. Pure Appl. Algebra* **219** (2015), no. 5, 1720–1741.
- [37] B. Olberding, Noetherian spaces of integrally closed rings with an application to intersections of valuation rings, *Comm. Algebra* **38** (2010), 3318–3332.
- [38] B. Olberding, Intersections of valuation overrings of two-dimensional Noetherian domains, in *Commutative algebra, Noetherian and non-Noetherian perspectives*, 335–361, Springer, New York, 2011.
- [39] B. Olberding, Overrings of two-dimensional Noetherian domains representable by Noetherian spaces of valuation rings *J. Pure Appl. Algebra* **212** (2008), no. 7, 1797–1821.
- [40] H. Uda, LCM-stableness in ring extensions, *Hiroshima Math. J.* **13** (1983), 357–377.
- [41] H. Weyl, Algebraic Theory of Numbers, Princeton University Press, Princeton, 1940.
- [42] O. Zariski and P. Samuel, Commutative Algebra, Volume II, Van Nostrand, Princeton, 1960.

E-mail address: `carmelo@mat.uniroma3.it`

E-mail address: `fontana@mat.uniroma3.it`

E-mail address: `spirito@mat.uniroma3.it`

DIPARTIMENTO DI MATEMATICA E FISICA, UNIVERSITÀ DEGLI STUDI “ROMA TRE”, ROMA, ITALY