

RING-THEORETIC PROPERTIES OF $PvMDS$

SAID EL BAGHDADI AND STEFANIA GABELLI

ABSTRACT. We extend to Prüfer v -multiplication domains some distinguished ring-theoretic properties of Prüfer domains. In particular we consider the $t\#\#$ -property, the t -radical trace property, w -divisoriality and w -stability.

INTRODUCTION

A Prüfer v -multiplication domain, for short a $PvMD$, is a domain whose localizations at t -maximal ideals are valuation domains [22]. For this reason, the ideal-theoretic properties of valuation domains globalize to t -ideals of $PvMDs$ and several properties of ideals of Prüfer domains hold for t -ideals of $PvMDs$: for example a domain is a $PvMD$ if and only if each t -finite t -ideal is t -invertible. The aim of this paper is to show that, introducing suitable t -analogues of some distinguished properties of integral domains, Prüfer domains and $PvMDs$ have a similar behaviour also from the ring-theoretic point of view. We recall that the class of $PvMDs$, besides Prüfer domains, includes Krull domains and GCD-domains.

As a matter of fact, the t -operation is not always as good as the w -operation for extending certain properties that hold in the classical case, that is in the d -operation setting. Thus in general it is often more convenient to consider the w -analogue of a given property (see for instance [43, 44, 9]). However in a $PvMD$ the w -operation and the t -operation coincide [30] and one can use indifferently these two star operations.

In Section 1, we deal with the $t\#\#$ -property and the $tRTP$ -property. $PvMDs$ satisfying the $t\#\#$ -property have been studied in [15, Section 2]. Here we characterize $PvMDs$ with the $tRTP$ -property; getting for example that a $PvMD$ is a $tRTP$ -domain if and only if each v -finite divisorial ideal has at most finitely many minimal primes. Then, generalizing the Prüfer case, we show that the $t\#\#$ -property and the $tRTP$ -property are strictly connected for $PvMDs$. Among other results, we prove that a $PvMD$ satisfying the $t\#\#$ -property is a $tRTP$ -domain and that the converse holds if each t -prime is branched. We also show that an almost Krull domain satisfying the $t\#\#$ -property is a Krull domain.

In Section 2 we introduce the notion of w -stability and relate it to w -divisoriality, a property defined and studied in [9]. First we show that the study of w -stability can be reduced to the t -local case. Then we use this result to extend to $PvMDs$ some properties of stable and divisorial Prüfer domains. For example, we prove that w -stability of t -primes enforces a $PvMD$ to be a generalized Krull domains and that an integrally closed w -stable domain is precisely a generalized Krull domain with t -finite character. We also characterize w -stable w -divisorial $PvMDs$ and show that these domains behave like totally divisorial Prüfer domains.

We assume that the reader is familiar with the language of star operations [18, Sections 32 and 34]. We recall some definitions and basic properties.

Date: June 12, 2006.

1991 Mathematics Subject Classification. Primary: 13F05; Secondary: 13A15, 13A18.

Key words and phrases. Prüfer v -multiplication domain, $\#$ -property, radical trace property, divisoriality, stability.

Throughout this paper R will denote an integral domain with quotient field K and we will assume that $R \neq K$.

A *star operation* is a map $I \rightarrow I^*$ from the set $F(R)$ of nonzero fractional ideals of R to itself such that:

- (1) $R^* = R$ and $(aI)^* = aI^*$, for all $a \in K \setminus \{0\}$;
- (2) $I \subseteq I^*$ and $I \subseteq J \Rightarrow I^* \subseteq J^*$;
- (3) $I^{**} = I^*$.

A star operation $*$ is of *finite type* if $I^* = \bigcup \{J^*; J \subseteq I \text{ and } J \text{ is finitely generated}\}$, for each $I \in F(R)$. To any star operation $*$, we can associate a star operation $*_f$ of finite type by defining $I^{*f} = \bigcup J^*$, with the union taken over all finitely generated ideals J contained in I . Clearly $I^{*f} \subseteq I^*$.

An ideal $I \in F(R)$ is a **-ideal* if $I = I^*$ and is **-finite* if $I^* = J^*$ for some finitely generated ideal J . A **-finite *-ideal* is also called a **-ideal of finite type*.

A **-prime* is a prime **-ideal* and a **-maximal* ideal is an ideal that is maximal in the set of the proper **-ideals*. A **-maximal* ideal (if it exists) is a prime ideal. If $*$ is a star operation of finite type, an easy application of Zorn's Lemma shows that the set $*\text{-Max}(R)$ of the **-maximal* ideals of R is not empty. In this case, for each $I \in F(R)$, $I^* = \bigcap_{M \in *\text{-Max}(R)} I^* R_M$; in particular $R = \bigcap_{M \in *\text{-Max}(R)} R_M$ [22].

For any star operation $*$, the set of **-ideals* of R is a semigroup under the **-multiplication*, defined by $(I, J) \mapsto (IJ)^*$, with unity R . An ideal $I \in F(R)$ is called **-invertible* if I^* is invertible with respect to the **-multiplication*. In this case the **-inverse* of I is $(R : I)$. Thus I is **-invertible* if and only if $(I(R : I))^* = R$.

The identity is a star operation, called the *d-operation*. The *v-operation* (or *divisorial closure*), the *t-operation* and the *w-operation* are the best known nontrivial star operations and are defined in the following way. For each $I \in F(R)$, we set $I_v := (R : (R : I))$ and $I_t := \bigcup J_v$ with the union taken over all finitely generated ideals J contained in I . Hence the *t-operation* is the finite type star operation associated to the *v-operation*. The *w-operation* is the star operation of finite type defined by setting $I_w := \bigcap_{M \in t\text{-Max}(R)} I R_M$. We have $w\text{-Max}(R) = t\text{-Max}(R)$ and $I R_M = I_w R_M \subseteq I_t R_M$, for each $M \in t\text{-Max}(R)$. Thus $I_w \subseteq I_t \subseteq I_v$. Also, an ideal $I \in F(R)$ is *w-invertible* if and only if it is *t-invertible*.

A Prüfer domain is an integrally closed domain such that $d = t$ [18, Proposition 34.12] and a PvMD is an integrally closed domain such that $w = t$ [30, Theorem 3.1].

The *v*-, *t*- and *w*-operations on R can be extended to the set of nonzero R -submodules of K by setting $E^v := (R : (R : E))$, $E_t = \bigcup \{F_v; F \subseteq E; F \text{ finitely generated}\}$ and $E_w = \bigcap \{E_M; M \in t\text{-Max}(R)\}$, for each non zero R -submodule E of K . In this way, one obtains *semistar operations* on R . For more details, see for example [12]. By viewing w as a semistar operation on R , we can say that an overring T of a domain R is *t-linked* over R if $T_w = T$ [4, Proposition 2.13]. Each overring of R is *t-linked* precisely when $d = w$ [4, Theorem 2.6].

We denote by $t\text{-Spec}(R)$ the set of *t*-prime ideals of R . Each height-one prime is a *t*-prime and each prime minimal over a *t*-ideal is a *t*-prime. We say that R has *t-dimension one* if each *t*-prime ideal has height one.

We now define the ring-theoretic properties considered in this paper.

The t###-property. The *#-property* and the *##-property* were introduced by R. Gilmer [19] and R. Gilmer and W. Heinzer [20] respectively. A domain R has the *#-property*, or it is a *#-domain*, if $\bigcap_{M \in \mathcal{M}_1} R_M \neq \bigcap_{M \in \mathcal{M}_2} R_M$, for any pair of distinct nonempty sets \mathcal{M}_1 and \mathcal{M}_2 of maximal ideals. Any overring of a one-dimensional Prüfer *#-domain* is a *#-domain* [19, Corollary 2], but in general

the $\#$ -property is not inherited by overrings [20, Section 2]. One says that R has the $\#\#$ -property, or it is a $\#\#$ -domain, if each overring of R is a $\#$ -domain.

The $t\#$ -property was introduced and studied in [15]. A domain R has the $t\#$ -property (or is a $t\#$ -domain) if $\bigcap_{M \in \mathcal{M}_1} R_M \neq \bigcap_{M \in \mathcal{M}_2} R_M$ for any two distinct subsets \mathcal{M}_1 and \mathcal{M}_2 of $t\text{-Max}(R)$. Although in [15] it was explored the transfer of the $t\#$ -property to some distinguished classes of overrings, it was not given any definition for the $t\#\#$ -property. Here, we say that R has the $t\#\#$ -property (or is a $t\#\#$ -domain) if $\bigcap_{P \in \mathcal{P}_1} R_P \neq \bigcap_{P \in \mathcal{P}_2} R_P$ for any two distinct subsets \mathcal{P}_1 and \mathcal{P}_2 of pairwise incomparable t -prime ideals of R . Our definition is motivated by the fact that for a PvMD this is equivalent to say that each t -linked overring has the $t\#$ -property [15, Proposition 2.10].

The tRTP-property. If R is an integral domain and M is a unitary R -module, the *trace* of M is the ideal of R generated by the set $\{f(m); f \in \text{Hom}_R(M, R), m \in M\}$. An ideal J of R is called a *trace ideal* or a *strong ideal* if it is the trace of some R -module M . This happens if and only if $J = I(R : I)$, for some nonzero ideal I of R , equivalently $(J : J) = (R : J)$ [10, Lemmas 4.2.2. and 4.2.3]. If V is a valuation domain, a trace ideal is either equal to V or it is prime [10, Proposition 4.2.1]; this fact led to the consideration of several conditions related to trace ideals [28]. The radical trace property was introduced by W. Heinzer and I. Papick [24] and is particularly significant for Prüfer domains [24, 33]. R is a domain satisfying the *radical trace property*, or it is an *RTP-domain*, if each proper strong ideal is a radical ideal, that is, for each nonzero ideal I of R , either $I(R : I) = R$ or $I(R : I)$ is a radical ideal.

A. Mimouni studied trace properties in the setting of star operations, in particular he considered the t -operation [35]. As in [35], we say that a domain R has the *t-radical trace property*, or it is a *tRTP-domain*, if each proper strong t -ideal of R is a radical ideal. This is equivalent to say that, for each nonzero ideal I of R , either $(I(R : I))_t = R$ or $(I(R : I))_t$ is a radical ideal.

w-divisoriability. The class of domains in which each nonzero ideal is divisorial has been studied, independently and with different methods, by H. Bass [2], E. Matlis [34] and W. Heinzer [23] in the sixties. Following S. Bazzoni and L. Salce [3], a domain in which each nonzero ideal is divisorial is now called a *divisorial domain* and a domain such that each overring is divisorial is called *totally divisorial*.

The most suitable star analogue of divisoriability is the notion of w -divisoriability that was introduced and extensively studied in [9]. A *w-divisorial domain* is a domain such that each w -ideal is divisorial.

w-stability. Motivated by earlier work of H. Bass [2] and J. Lipman [32] on the number of generators of an ideal, in 1974 J. Sally and W. Vasconcelos defined a Noetherian ring R to be *stable* if each nonzero ideal of R is projective over its endomorphism ring $\text{End}_R(I)$ [41, 42]. When I is a nonzero ideal of a domain R , then $\text{End}_R(I) = (I : I)$; thus a domain R is stable if each nonzero ideal I of R is invertible in the overring $(I : I)$. B. Olberding showed that stability and divisoriability are strictly connected and that stability is particularly significant in the context of Prüfer domains [10, 36, 37, 38, 39].

We introduce the notion of w -stability in Section 2. We say that a w -ideal I of a domain R is *w-stable* if I is w -invertible in the overring $E(I) := (I : I)$, that is $(I(E(I) : I))_w = E(I)$, and say that R is *w-stable* if each w -ideal of R is w -stable. For a more general notion of stability with respect to a semistar operation we refer the reader to the forthcoming paper [16].

1. $t\#\#$ -PROPERTY AND t -RADICAL TRACE PROPERTY

The $\#\#$ -property and the radical trace property are closely related for a Prüfer domain. In this section we compare the t -analogues of these two properties for PvMDs.

Several characterizations of PvMDs satisfying the $t\#\#$ -property have been given in [15, Section 2]. For the study of the $tRTP$ -property, we need some results on branched t -primes. Recall that a prime ideal P of a domain R is *branched* if there exists a P -primary ideal distinct from P . Clearly P is branched if and only if PR_P is branched. Since the localization of a PvMD at a t -prime is a valuation domain, the branched t -primes of PvMDs can be characterized by properties similar to those well known for the branched primes of Prüfer domains [18, Theorem 23.3 (e)].

Lemma 1.1. *Let R be a PvMD and $J := x_1R + \cdots + x_nR$ a nonzero finitely generated ideal such that $J_v \neq R$. If P is a t -prime containing J , then P is minimal over J_v if and only if P is minimal over J , if and only if P is minimal over x_iR , for some i , $1 \leq i \leq n$.*

Proof. It is enough to observe that, since R_P is a valuation domain, we have $J_vR_P = J_tR_P = JR_P = x_iR_P$, for some i , $1 \leq i \leq n$. \square

Proposition 1.2. *Let R be a PvMD and P a t -prime of R . The following conditions are equivalent:*

- (i) P is branched;
- (ii) P is a minimal prime of a principal ideal;
- (iii) P is a minimal prime of a finitely generated ideal;
- (iv) P is a minimal prime of a v -finite divisorial ideal;
- (v) P is not the union of the set of (t) -primes of R properly contained in P .

Proof. The equivalence of conditions (i), (ii) and (v) is obtained by localizing at P and using [18, Theorem 17.3 (e)]. The equivalence of conditions (ii), (iii) and (iv) follows from Lemma 1.1. \square

In any commutative ring with unity, if each minimal prime of an ideal I is the radical of a finitely generated ideal, then I has only finitely many minimal primes [21, Theorem 1.6]. By passing through the t -Nagata ring, we now show that a similar result holds for t -ideals of PvMDs.

If R is an integral domain, we set $R(X) := R[X]_N$, where $N = \{f \in R[X] : c(f) = R\}$ and $R\langle X \rangle := R[X]_{N_t}$, where $N_t = \{f \in R[X] : c(f)_t = R\}$. $R(X)$ is called the *Nagata ring* of R and $R\langle X \rangle$ the *t -Nagata ring* of R [12, 29, 30].

B. G. Kang proved that R is a PvMD if and only if $R\langle X \rangle$ is a Prüfer (indeed a Bezout) domain [30, Theorem 3.7]. In addition, there is a lattice isomorphism between the lattice of t -ideals of R and the lattice of ideals of $R\langle X \rangle$ [30, Theorem 3.4]. More precisely, we have:

Proposition 1.3. *Let R be a PvMD. Then the map $I_t \mapsto IR\langle X \rangle$ is an order-preserving bijection between the set of t -ideals of R and the set of nonzero ideals of $R\langle X \rangle$, whose inverse is the map $J \mapsto J \cap R$. Moreover, P is a t -prime (respectively, t -maximal) ideal of R if and only if $PR\langle X \rangle$ is a prime (respectively, maximal) ideal of $R\langle X \rangle$ and we have $R\langle X \rangle_{PR\langle X \rangle} = R[X]_{PR[X]} = R_P(X)$.*

Proposition 1.4. *Let R be a PvMD and I a proper t -ideal of R . If each minimal prime of I is the radical of a v -finite divisorial ideal, then I has finitely many minimal t -primes.*

Proof. Each minimal prime of a t -ideal I is a t -ideal of R . By Proposition 1.3, the map $P \mapsto PR\langle X \rangle$ is a bijection between the set of minimal primes of I and the set of minimal primes of $IR\langle X \rangle$. Moreover if J is a nonzero finitely generated ideal of R such that $P = \text{rad}(J_v)$, then $PR\langle X \rangle = \text{rad}(JR\langle X \rangle)$. Hence each minimal prime of $IR\langle X \rangle$ is the radical of a finitely generated ideal. By [21, Theorem 1.6], $IR\langle X \rangle$ has finitely many minimal primes and the same holds for I . \square

If T is an overring of R , the w -operation and the t -operation on R , viewed as semistar operations, induce two semistar operations of finite type on T , which here are still denoted by w and t respectively. If in addition T is t -linked over R , the w -operation is a star operation on T [8, Proposition 3.16]. Note that this star operation, being spectral and of finite type [12], is generally smaller than the w -operation on T , that we denote by w' to avoid confusion.

Proposition 1.5. *Let R be a PvMD and T a t -linked overring of R . Then T is a PvMD and $w = t = t' = w'$ on T , where w' and t' denote respectively the w -operation and the t -operation on T .*

Proof. When R is a PvMD also T is a PvMD [30, Theorem 3.8 and Corollary 3.9]. In addition, if R is a PvMD the two semistar operations w and t coincide [11, Theorem 3.1 ((i) \Rightarrow (vi))]. Hence $w = t$ and $w' = t'$ as star operations on T .

We next show that $t = t'$ on T . Let I be a nonzero ideal of T . Clearly, $I_t \subseteq I_{t'}$. On the other hand, we have $I_{t'} = \bigcap \{IT_M; M \in t'\text{-Max}(T)\}$. Since $I_t = \bigcap \{I_t T_N; N \in t\text{-Max}(T)\}$ [22, Proposition 4], to show that $I_{t'} \subseteq I_t$ it suffices to show that $t\text{-Max}(T) \subseteq t'\text{-Max}(T)$.

If $N \in t\text{-Max}(T)$, we have that $(N \cap R)_t \subseteq N_t \cap R_t = N \cap R$. Hence $N \cap R$ is a t -prime of R and $T_N \supseteq R_{N \cap R}$ are valuation domains. It follows that N is a t' -prime of T . In addition N is t' -maximal because it is t -maximal and each t' -prime of T is also a t -prime. \square

If I is a w -ideal of R , then it is easily shown that $E(I)_w = E(I)$. Thus $E(I)$ is a t -linked overring of R . It follows that, when R is a PvMD, by Proposition 1.5, $E(I)$ is a PvMD and $w = t = t' = w'$ on $E(I)$.

Proposition 1.6. *Let R be a PvMD. Then:*

- (1) *If I is a strong t -ideal of R , then $E(I) = S \cap T$, where $S = \bigcap_{P \in \text{Min}(I)} R_P$ and $T := \bigcap_{M \in t\text{-Max}(R), M \not\supseteq I} R_M$.*
- (2) *If P is a t -prime of R which is not t -invertible, then $E(P) = (R : P) = R_P \cap T$, where $T := \bigcap_{M \in t\text{-Max}(R), M \not\supseteq P} R_M$.*
- (3) *If P is a t -prime of R which is not t -invertible and R satisfies the ascending chain condition on radical t -ideals, then P is t -maximal in $E(P)$.*

Proof. (1) follows from [27, Theorem 4.5].

(2) If P is not t -invertible, then P is strong by [26, Proposition 2.3 and Lemma 1.2]. Whence (2) follows from (1).

(3) By part (2), $E(P) = (R : P)$. Since $E(P)$ is t -linked over R , then $E(P)$ is t -flat on R (that is $E(P)_Q = R_{Q \cap R}$ for each t -prime ideal Q of $E(P)$) [31, Proposition 2.10] and P is a t -ideal of $E(P)$ (Proposition 1.5). Let Q be a t -prime of $E(P)$ properly containing P . By t -flatness we can write $Q = (P'E(P))_t$, where $P' = Q \cap R$ is a t -prime of R properly containing P [6, Proposition 2.4]. By the ascending chain condition on radical t -ideals, $P' = \text{rad}(J_t)$ for some finitely generated ideal J [6, Lemma 3.7]. Since $P \subsetneq P'$, by checking t -locally, we get that $P \subsetneq J_t$. We have $R = (J(R : J))_t \subseteq (J(R : P))_t = (JE(P))_t \subseteq (P'E(P))_t = Q$. A contradiction. Hence $P = Q$ and so P is a t -maximal ideal of $E(P)$. \square

Lemma 1.7. *Let R be a PvMD satisfying the t RTP-property. If P is a branched t -prime of R which is not t -invertible, then $R_P \not\supseteq T$, where $T = \bigcap_{M \in t\text{-Max}(R), M \not\supseteq P} R_M$.*

Proof. If $R_P \supseteq T$ then, by Proposition 1.6 (2), $T = E(P)$. Let Q be a P -primary ideal of R . Since P is a t -ideal, we may assume that Q is a t -ideal. We have $QT \subseteq PT = P \subseteq R$ and so $QT \subseteq QR_P \cap R = Q$. Hence $QT = Q$. If M is a t -maximal ideal of R such that $P \not\subseteq M$, then $Q \not\subseteq M$. Thus $(R : Q) \subseteq R_M$ and it follows that $(R : Q) \subseteq T$. Hence $Q(R : Q) = Q$. Since R is a t RTP-domain, then we must have $Q = P$. It follows that P is not branched. \square

Theorem 1.8. *Let R be a PvMD. The following conditions are equivalent:*

- (i) *R is a t RTP-domain;*

- (ii) Each branched t -prime P contains a finitely generated ideal J such that $J \subseteq P$ and $J \not\subseteq M$, for each $M \in t\text{-Max}(R)$ not containing P ;
- (iii) Each branched t -prime is the radical of a v -finite divisorial ideal;
- (iv) Each nonzero principal ideal has at most finitely many minimal (t) -primes;
- (v) Each nonzero finitely generated ideal has at most finitely many minimal t -primes;
- (vi) Each v -finite divisorial ideal has at most finitely many minimal (t) -primes.

Proof. (i) \Rightarrow (ii). Let P be a branched t -prime of R . If P is t -invertible, then P is v -finite and there is nothing to prove. If P is not t -invertible, then $E(P) = (R : P) = R_P \cap T$, where $T = \bigcap_{M \in t\text{-Max}(R), M \not\supseteq P} R_M$ (Proposition 1.6 (2)). Since $R_P \not\subseteq T$ (Lemma 1.7), there exists a finitely generated ideal J such that $J \subseteq P$ and $J \not\subseteq M$, for each $M \in t\text{-Max}(R)$ not containing P [7, Lemma 3.6].

(ii) \Rightarrow (iii) Let P be a branched t -prime of R and J as in the hypothesis. By Proposition 1.2, P is minimal over a finitely generated ideal H . Hence P is the radical of the v -finite divisorial ideal $(J + H)_v$.

(iii) \Rightarrow (iv). Let $x \in R$ be a nonzero nonunit and let $\{P_\alpha\}$ be the set of minimal primes of xR . By Proposition 1.2 each P_α is branched. Hence by hypothesis each P_α is the radical of a v -finite divisorial ideal. It follows from Proposition 1.4 that $\{P_\alpha\}$ is a finite set.

(iv) \Leftrightarrow (v) \Leftrightarrow (vi) follow from Lemma 1.1.

(iv) \Rightarrow (iii). Let P be a branched t -prime. By Proposition 1.2, P is minimal over a v -finite divisorial ideal I . Since I has finitely many minimal primes, then P is the radical of a v -finite divisorial ideal by [15, Lemma 2.13].

(iii) \Rightarrow (i). By [35, Theorem 15] it is enough to show that for each strong t -ideal I and each minimal prime P of I we have $IR_P = PR_P$.

Assume that $IR_P \subsetneq PR_P$. Then P is branched, because R_P is a valuation domain. Thus $P = \text{rad}(H_v)$ for some finitely generated ideal H . Let $a \in P$ be such that $IR_P \subsetneq aR_P \subseteq PR_P$ and set $J = H + aR$. Then P is the radical of J_v . By checking t -locally, we have that $I \subseteq J_v$. In fact, let $M \in t\text{-Max}(R)$. If $P \not\subseteq M$, then $IR_M \subseteq R_M = J_v R_M$. If $P \subseteq M$, then $IR_M \subseteq IR_P$. Hence $a \notin IR_M$ and $IR_M \subsetneq J_v R_M$. Since I is strong, by Proposition 1.6(1), $(R : I) = E(I) \subseteq R_P$. Whence $J(R : J) \subseteq P(R : I) \subseteq PR_P$ and so $J(R : J) \subseteq P$. A contradiction because J is t -invertible. \square

Since in a Prüfer domain the t -operation is trivial, we get the following corollary, due to T. Lucas. The equivalence (i) \Leftrightarrow (ii) is [33, Theorem 23], while (i) \Leftrightarrow (iv) is, to our knowledge, unpublished.

Corollary 1.9. *Let R be a Prüfer domain. The following conditions are equivalent:*

- (i) R is an RTP-domain;
- (ii) Each branched prime is the radical of a finitely generated ideal;
- (iii) Each principal ideal has at most finitely many minimal primes;
- (iv) Each finitely generated ideal has at most finitely many minimal primes.

The following theorem was stated for Prüfer domains in [33, Corollaries 25 and 26].

Theorem 1.10. *Let R be a PvMD.*

- (1) If R is a $t\#\#\text{-domain}$, then R is a $t\text{RTP}$ -domain.
- (2) If R is a $t\text{RTP}$ -domain and each t -prime is branched, then R is a $t\#\#\text{-domain}$.

Proof. A PvMD R has the $t\#\#\text{-property}$ if and only if, for each t -prime ideal P , there exists a finitely generated ideal $J \subseteq P$ such that each t -maximal ideal containing J must contain P [15, Proposition 2.8]. Hence we can apply Theorem 1.8, (i) \Leftrightarrow (ii). \square

Theorem 1.11. *Let R be a PvMD. The following conditions are equivalent:*

- (i) R satisfies the ascending chain condition on radical t -ideals;
- (ii) R is a $tRTP$ -domain satisfying the ascending chain condition on prime t -ideals;
- (iii) R is a $t\#\#\text{-domain}$ satisfying the ascending chain condition on prime t -ideals;
- (iv) R is a $t\#\#\text{-domain}$ and each t -prime is branched.

Proof. (i) \Leftrightarrow (iii) \Leftrightarrow (iv) by [15, Proposition 2.14].

(ii) \Leftrightarrow (iii) By Proposition 1.2, the ascending chain condition on prime t -ideals implies that each t -prime of R is branched. Hence we can apply Theorem 1.10. \square

Recall that a domain R has *finite character* (respectively, *t -finite character*) if each nonzero element of R belongs to at most finitely many maximal (respectively, t -maximal) ideals. A domain with finite character such that each nonzero prime ideal is contained in a unique maximal ideal was called by E. Matlis an *h -local domain*. Following [1], we say that R is a *weakly Matlis domain* if R has t -finite character and each t -prime ideal is contained in a unique t -maximal ideal.

Theorem 1.12. *Let R be a PvMD and consider the following conditions:*

- (i) R is a weakly Matlis domain;
- (ii) R has t -finite character;
- (iii) R has the $t\#\#\text{-property}$;
- (iv) R is a $tRTP$ -domain.

Then (i) \Rightarrow (ii) \Rightarrow (iii) \Rightarrow (iv).

If in addition each t -prime ideal of R is contained in a unique t -maximal ideal, all these conditions are equivalent.

Proof. (i) \Rightarrow (ii) by definition.

(ii) \Rightarrow (iii). If R has t -finite character, for each $\Lambda \subseteq t\text{-Max}(R)$ the multiplicative system of ideals $\mathcal{F}(\Lambda) := \{I; I \not\subseteq M, \text{ for each } M \in \Lambda\}$ is finitely generated [14, Proposition 2.7]. We conclude that R is a $t\#\#\text{-domain}$ by applying [15, Proposition 2.8].

(iii) \Rightarrow (iv) By Theorem 1.10 (1).

Now assume that each t -prime of R is contained in a unique t -maximal ideal. Then clearly conditions (i) and (ii) are equivalent.

(iv) \Rightarrow (ii) By Theorem 1.8, for each nonzero nonunit $x \in R$, the ideal xR has finitely many minimal (t)-primes. Since each t -prime is contained in a unique t -maximal ideal, then x is contained in finitely many t -maximal ideals. \square

When R is a Prüfer domain, for $d = t$, from Theorem 1.11 we get [24, Theorem 2.7] and from Theorem 1.12 we get [36, Proposition 3.4].

Remark 1.13. The hypothesis that R is a PvMD in Theorems 1.11 and 1.12 cannot be relaxed. In fact each Noetherian domain is a $t\#\#\text{-domain}$ [15, Proposition 2.4], but it is not necessarily a $tRTP$ -domain [24, Corollary 2.2].

A *strongly discrete valuation domain* is a valuation domain such that each nonzero prime ideal is not idempotent [10, p. 145] and a *strongly discrete Prüfer domain* is a domain whose localizations at nonzero prime ideals are strongly discrete valuation domains; equivalently a domain such that $P \neq P^2$ for each nonzero prime ideal P [10, Proposition 5.3.5]. We say that a PvMD R is *strongly discrete* if R_P is a strongly discrete valuation domain for each t -prime ideal P of R ; equivalently, if $(P^2)_t \neq P$, for each $P \in t\text{-Spec}(R)$ [9, Lemma 3.4]. *Generalized Krull domains* were introduced by the first author in [6] and can be defined as strongly discrete PvMDs satisfying the ascending

chain condition on radical t -ideals [6, Theorem 3.5 and Lemma 3.7]. In the Prüfer case, that is for $d = t$, this class of domains coincides with the class of *generalized Dedekind domains* introduced by N. Popescu in [40]. A Krull domain is a generalized Krull domain of t -dimension one [6, Theorem 3.11].

Theorem 1.14. *Let R be an integral domain. The following conditions are equivalent:*

- (i) R is a generalized Krull domain;
- (ii) R is a strongly discrete PvMD satisfying the $t\#\#$ -property;
- (iii) R is a strongly discrete PvMD satisfying the $tRTP$ -property.

Proof. (i) \Rightarrow (ii) follows from Theorem 1.11 and (ii) \Rightarrow (iii) follows from Theorem 1.10(1).

(iii) \Rightarrow (i). By Theorem 1.8, R is a strongly discrete PvMD such that each proper v -finite divisorial ideal has finitely many minimal primes. Hence R is a generalized Krull domain by [6, Theorem 3.9]. \square

In the Prüfer case, for $d = t$, we recover from Theorem 1.14 a well known characterization of generalized Dedekind domains, see for example [10, Theorem 5.5.4].

Remark 1.15. Any w -divisorial domain is a $t\#$ -domain. In fact, clearly all the t -maximal ideals of a w -divisorial domain are divisorial; hence we can apply [15, Theorem 1.2].

If R is a domain such that $R_{\mathcal{F}(\Lambda)} := \bigcap_{P \in \Lambda} R_P$ is w -divisorial, for each set Λ of pairwise incomparable t -primes, then $\mathcal{F}(\Lambda)$ is v -finite by [9, Proposition 2.2]. Thus $t\text{-Max}(R_{\mathcal{F}(\Lambda)}) = \{P_{\mathcal{F}(\Lambda)}; P \in \Lambda\}$ [9, Lemma 2.1]. It follows that, given two different sets Λ_1 and Λ_2 of pairwise incomparable t -primes of R , we have $R_{\mathcal{F}(\Lambda_1)} \neq R_{\mathcal{F}(\Lambda_2)}$. Therefore R is a $t\#\#$ -domain.

Conversely, it is not true that a $t\#\#$ -domain is w -divisorial. In fact each Noetherian domain has the $t\#\#$ -property [15, Proposition 2.4], but a w -divisorial Noetherian domain must have t -dimension one [9, Theorem 4.2].

An integral domain R is an *almost Krull domain* if R_M is a rank-one discrete valuation domain for each t -maximal ideal M of R . Almost Krull domains were studied by Kang under the name of t -almost Dedekind domains in [30, Section IV]. A Krull domain is an almost Krull domain with t -finite character. In dimension one, the class of almost Krull domains coincides with the class of almost Dedekind domains introduced by R. Gilmer [17]. Gilmer showed that an almost Dedekind domain satisfying the $\#$ -property must be Dedekind [19, Theorem 3]. Next, we extend this result to almost Krull domains. First, we give the following characterization of almost Krull domains, which follows directly from the definitions.

Lemma 1.16. *Let R be an integral domain. Then R is an almost Krull domain if and only if R is a strongly discrete PvMD of t -dimension one.*

Theorem 1.17. *Let R be an integral domain. Then the following conditions are equivalent:*

- (i) R is an almost Krull domain satisfying the $t\#$ -property;
- (ii) R is an almost Krull domain satisfying the $t\#\#$ -property;
- (iii) R is a Krull domain.

Proof. (i) \Rightarrow (ii) Since an almost Krull domain has t -dimension one (Lemma 1.16).

(ii) \Rightarrow (iii) By Lemma 1.16 and Theorem 1.14, if (ii) holds, R is a generalized Krull domain of t -dimension 1. Hence R is a Krull domain by [6, Theorem 3.11].

(iii) \Rightarrow (i) Follows from Theorem 1.14. \square

We end this Section by putting into evidence that the $t\#\#$ -property and the $tRTP$ -property of a PvMD are related respectively to the $\#\#$ -property and the RTP -property of its t -Nagata ring.

Theorem 1.18. *Let R be a PvMD. Then:*

- (1) *R is a $t\#$ -domain if and only if $R\langle X \rangle$ is a $\#$ -domain.*
- (2) *R is a $t\#\#$ -domain if and only if $R\langle X \rangle$ is a $\#\#$ -domain.*
- (3) *R is a $tRTP$ -domain if and only if $R\langle X \rangle$ is a RTP -domain.*

Proof. (1) follows from [15, Theorem 3.6]. The proof of (2) is similar and it is obtained by using Proposition 1.3 and the characterization of Prüfer $\#\#$ -domains and PvMDs satisfying the $t\#\#$ -property proved respectively in [20, Theorem 3] and [15, Proposition 2.8 (7)].

(3) Follows from Proposition 1.3, Theorem 1.8 (v) and Corollary 1.9 (iv). \square

2. w -DIVISORIALITY AND w -STABILITY

The notion of w -divisoriality has been studied in [9]. A domain R is an integrally closed w -divisorial domain if and only if it is a weakly Matlis PvMD such that each t -maximal ideal is t -invertible [9, Theorem 3.3] and R is an integrally closed domain such that each t -linked overring is w -divisorial if and only if it is a weakly Matlis strongly discrete PvMD, equivalently R is a w -divisorial generalized Krull domain [9, Theorem 3.5]. We now introduce the notion of w -stability and show that in PvMDs w -divisoriality and w -stability are strictly related; thus extending some results proved by B. Olberding for Prüfer domains.

As before, if T is a t -linked overring of R , we denote by w the star operation induced on T by the w -operation on R and by w' the w -operation on T . We say that a w -ideal I of R is w -stable if I is w -invertible in the (t -linked) overring $E(I) := (I : I)$, that is if $(I(E(I) : I))_w = E(I)$, and we say that R is w -stable if each w -ideal of R is w -stable.

Our first result is a generalization of [39, Theorems 3.3 and 3.5] and shows in particular that the study of w -divisorial domains can be reduced to the t -local case. We recall that a valuation domain is stable if and only if it is strongly discrete [10, Proposition 5.3.8].

Lemma 2.1. *Let R be a quasi-local domain. Then a nonzero ideal I of R is stable if and only if $I^2 = xI$ for some $x \in I$.*

Proof. This follows from [39, Lemma 3.1] and [10, Lemma 7.3.4]. \square

Theorem 2.2. *Let R be an integral domain. The following conditions are equivalent:*

- (i) *R is w -stable;*
- (ii) *Each w -ideal I of R is divisorial in $E(I)$;*
- (iii) *R has t -finite character and R_M is stable for each t -maximal ideal M of R .*

Proof. (i) \Rightarrow (ii). Let I be a w -ideal of R . Denote by v' the v -operation on $E(I)$. Since I is w -stable, we have $E(I) = (I(E(I) : I))_w \subseteq (I_{v'}(E(I) : I))_w \subseteq E(I)$. Hence $(I_{v'}(E(I) : I))_w = E(I)$ and $I = IE(I) = (I(E(I) : I)I_{v'})_w = ((I(E(I) : I))_w I_{v'})_w = I_{v'}$.

(ii) \Rightarrow (i). Let I be a w -ideal of R and set $J = (E(I) : I)$. Proceeding like in the proof of [39, Theorem 3.5 ((2) \Rightarrow (1))], we have $(E(I) : IJ) = E(I)$ and hence $E((IJ)_w) = E(I)$. Thus $(IJ)_w$ is a divisorial ideal of $E(I)$. It follows that $(IJ)_w = (E(I) : (E(I) : IJ)) = E(I) : E(I) = E(I)$, that is I is a w -stable ideal.

(i) \Rightarrow (iii) Let M be a t -maximal ideal of R and let $I = JR_M$ be a nonzero ideal of R_M , where J is an ideal of R which can be assumed to be a w -ideal (since $J_w R_M = JR_M$). By w -stability in R , $(J(E(J) : J))_w = E(J)$; in particular, $J(E(J) : J)R_M = E(J)R_M$. Since $1 \in E(J)R_M = J(E(J) : J)R_M \subseteq I(E(I) : I) \subseteq E(I)$, then $I(E(I) : I) = E(I)$. Hence I is a stable ideal of R_M and therefore R_M is stable.

We next show that R has t -finite character. Let M be a t -maximal ideal of R . Since R_M is a quasi-local stable domain, then $M^2 R_M = m M R_M$ for some $m \in M$ (Lemma 2.1). Set $I(M) :=$

$mR_M \cap R$. The ideal $I(M)$ is a t -ideal of R and $M^2 \subseteq I(M)$. Hence M is the only t -maximal ideal of R containing $I(M)$. From this, and by checking t -locally, we get $(I(M) : I(M)) = R$. Since R is w -stable, $I(M)$ is a w -invertible ideal of R . Thus $I(M)$ is divisorial.

Now, let $\{M_\alpha\}$ be a family of t -maximal ideals of R such that $\bigcap M_\alpha \neq (0)$. We want to show that $\{M_\alpha\}$ is a finite family.

Set $I_\alpha := I(M_\alpha)$ and let $J_\alpha := (\sum_{\beta \neq \alpha} (R : I_\beta))_w$. Note that J_α is a fractional ideal of R since $\bigcap_{\beta \neq \alpha} I_\beta \supseteq \bigcap_{\beta \neq \alpha} M_\beta^2 \neq 0$. We claim that $(J_\alpha : J_\alpha) = R$. To show this, we prove that $(J_\alpha : J_\alpha) \subseteq R_M$ for each t -maximal ideal M of R . Let $x \in (J_\alpha : J_\alpha)$. We first assume that $M \notin \{M_\beta\}_{\beta \neq \alpha}$. Since M_β is the only t -maximal ideal of R containing I_β , then $(R : I_\beta) \subseteq R_M$ for each $\beta \neq \alpha$. Hence $x \in xJ_\alpha \subseteq J_\alpha \subseteq R_M$. If $M = M_\gamma$ for some $\gamma \neq \alpha$. We have $x(R : I_\gamma) \subseteq (\sum_{\beta \neq \alpha} (R : I_\beta))_w$, and since I_γ is w -invertible, then $x \in (\sum_{\beta \neq \alpha} (I_\gamma(R : I_\beta))_w)_w$. Moreover, for $\beta = \gamma$, we have $(I_\gamma(R : I_\gamma))_w = R$, and for $\beta \neq \gamma$, $(R : I_\beta) \subseteq R_{M_\gamma}$. Hence $(\sum_{\beta \neq \alpha} (I_\gamma(R : I_\beta))_w)_w \subseteq R_{M_\gamma}$. Thus $x \in R_M$, which prove the claim.

Now, for each α , set $T_\alpha := \bigcap_{\beta \neq \alpha} M_\beta$. We claim that $T_\alpha \not\subseteq N$ for each t -maximal ideal $N \notin \{M_\beta\}_{\beta \neq \alpha}$. By the w -stability, J_α is a w -invertible ideal of $(J_\alpha : J_\alpha) = R$. In particular, J_α is w -finite. Thus $(R : J_\alpha)_N = (R_N : J_{\alpha N}) = R_N$ (since $I_\beta \not\subseteq N$ for each $\beta \neq \alpha$). On the other hand, we have $(R : J_\alpha)_N = (R : \sum_{\beta \neq \alpha} (R : I_\beta))R_N = (\bigcap_{\beta \neq \alpha} I_\beta)R_N$. Thus $\bigcap_{\beta \neq \alpha} I_\beta \not\subseteq N$. Since $\bigcap_{\beta \neq \alpha} I_\beta \subseteq T_\alpha$, then $T_\alpha \not\subseteq N$, in particular, $T_\alpha \not\subseteq M_\alpha$ for each α .

Now we proceed as in the proof of [25, Theorem 3.1]. Set $T := \sum T_\alpha$. By the above result T is not contained in any t -maximal ideal of R , hence $T_t = R$. Thus $(\sum_{i=1}^n T_i)_t = R$ for some finite subset $\{T_1, \dots, T_n\}$ of $\{T_\alpha\}$. Let $\{M_1, \dots, M_n\}$ be the corresponding set of t -maximal ideals. If $M_\alpha \notin \{M_1, \dots, M_n\}$ for some α , then $\sum_{i=1}^n T_i \subseteq M_\alpha$, which is impossible. Hence $\{M_\alpha\}$ is finite. Therefore R has t -finite character.

(iii) \Rightarrow (i) Let I be a w -ideal of R and let M_1, \dots, M_n be the t -maximal ideals of R containing I . Since I is t -locally stable then $IR_{M_i} = J_i E(I_{M_i})$ for some finitely generated ideal $J_i \subseteq I$, $i = 1, \dots, n$. Choose $y \in I$ such that $y \notin M$ for each t -maximal ideal $M \neq M_i$ containing the ideal $H := \sum J_i$ and consider the ideal $J := H + Ry$ of R . Clearly J is finitely generated. One can easily check that $IR_N = JE(I_N)$ for each t -maximal ideal N of R . We next show that $E(I_N) = E(I)_N$ for each t -maximal ideal N of R . Let $x \in E(I_N)$. Since $I_N = JE(I_N)$, then $xJ \subseteq I_N$. Hence $sxJ \subseteq I$ for some $s \in R \setminus N$. Let M be a t -maximal ideal of R . Then $sxI_M = sxJE(I_M) \subseteq IE(I_M) \subseteq I_M$. Thus $sxI_M \subseteq I_M$ for each t -maximal ideal M of R , so that $sxI \subseteq I$. Hence $x \in E(I)_N$. It follows that $E(I_N) = E(I)_N$, for each t -maximal ideal N , and $I = \bigcap_N I_N = \bigcap_N JE(I_N) = \bigcap_N JE(I)_N = (JE(I))_w$.

Finally, $(I(E(I) : I))_N = I_N(E(I) : JE(I))_N = I_N(E(I)_N : JE(I)_N) = I_N(E(I_N) : I_N) = E(I_N) = E(I)_N$, for each t -maximal ideal N . Therefore $(I(E(I) : I))_w = E(I)$ and so I is a w -stable ideal of R . \square

Proposition 2.3. *Let R be a w -stable domain. Then:*

- (1) *Each t -maximal ideal of R is divisorial.*
- (2) *t -Spec(R) is treed.*
- (3) *R satisfies the ascending chain condition on prime t -ideals.*

Proof. (1) Let M be a t -maximal ideal of R . If M is not divisorial, then $M_v = R$. Thus $E(M) = (R : M) = R$ and M is t -invertible. Hence M is divisorial, which is impossible.

(2) and (3) follow from Theorem 2.2 because a quasi-local stable domains has these properties [39, Theorem 4.11]. \square

The previous proposition shows that w -stable domains have some properties in common with generalized Krull domains [6]. We now prove that w -stability of t -primes enforces a PvMD to be a generalized Krull domain. For Prüfer domains, this follows from [13, Theorem 5] or [36, Theorem 4.7].

Theorem 2.4. *Let R be a PvMD. The following conditions are equivalent:*

- (i) R is a generalized Krull domain;
- (ii) Each radical t -ideal of R is divisorial and each divisorial ideal is w -stable;
- (iii) Each radical t -ideal of R is w -stable;
- (iv) Each t -prime ideal of R is w -stable.

Proof. We shall freely use Proposition 1.5.

(i) \Rightarrow (ii). Since $t\text{-Spec}(R)$ is treed and a t -ideal of R has finitely many minimal primes, a radical t -ideal of R is a t -product of finitely many t -primes [7, Lemma 2.5]. Hence each radical t -ideal of R is divisorial [7, Proposition 3.1].

Let I be a divisorial ideal of R . If I is t -invertible, hence w -invertible, then $E(I) = R$ and so I is w -stable. If I is not t -invertible, then consider the ideal $H := (I(R : I))_w$. By [7, Proposition 2.6], we have $H = (P_1 \cdots P_n)_w$, where $n \geq 1$ and each P_i is a strong t -prime. Thus $E(P_i) = (R : P_i) \subseteq (R : H) = (R : I(R : I)) = ((R : (R : I)) : I) = E(I)$. Since P_i is t -maximal in $E(P_i)$ (Proposition 1.6) and $E(P_i)$ is t -linked over R then P_i is t -invertible in $E(P_i)$ by [6, Corollaries 3.2 and 3.6]. Hence $(P_i(E(P_i) : P_i))_w = (P_i(E(P_i) : P_i))_t = (P_i(E(P_i) : P_i))_{t'} = E(P_i)$, where t' denotes the t -operation on $E(P_i)$. Thus $P_i E(I)$ is w -invertible in $E(I)$, for each i . It follows that H is w -invertible in $E(I)$ and so I has the same property.

(ii) \Rightarrow (iii) \Rightarrow (iv) are clear.

(iv) \Rightarrow (i). Let P be a t -prime of R . Since P is w -invertible in $E(P)$, then $P \neq (P^2)_w$ and so $P \neq (P^2)_t$. Thus R is a strongly discrete PvMD.

To prove that R is a generalized Krull domain, it is enough to show that R has the $t\#\#$ -property (Theorem 1.14). Let T be a t -linked overring of R and denote by t' the t -operation on T . Let M be a t' -maximal ideal of T . Since T is a PvMD, then $T = E(M)$. The ideal $P = M \cap R$ is a t -prime of R and $M = (PT)_{t'} = (PT)_w$ (cf. [31, Proposition 2.10] and [6, Proposition 2.4]). Thus $R \subseteq E(P) \subseteq E(M) = T$. Since P is w -stable, then PT is w -invertible in T , and hence M is w -invertible in T . So, M is a t' -invertible t' -ideal of T (since $w = t'$ in T). In particular M is a divisorial ideal of T . We conclude that T is a $t\#\#$ -domain by applying [15, Theorem 1.2]. \square

It is known that a generalized Dedekind domain need not be stable [13, Example 10]. In fact an integrally closed domain is stable if and only if it is a strongly discrete Prüfer domain with finite character [36, Theorem 4.6]. Hence a generalized Dedekind domain is stable if and only if it has finite character. We now extend these results to generalized Krull domains.

Lemma 2.5. *A domain with t -finite character is a strongly discrete PvMD if and only if it is a generalized Krull domain.*

Proof. A strongly discrete PvMD is a generalized Krull domain if and only if each nonzero nonunit has finitely many minimal t -primes [6, Theorem 3.9]. We conclude by recalling that in a PvMD two incomparable t -primes are t -comaximal. \square

Theorem 2.6. *Let R be an integral domain. The following conditions are equivalent:*

- (i) R is integrally closed and w -stable;
- (ii) R is a w -stable PvMD;
- (iii) R is a strongly discrete PvMD with t -finite character;

- (iv) R is a generalized Krull domain with t -finite character;
- (v) R is a w -stable generalized Krull domain;
- (vi) R is a PvMD with t -finite character and each t -prime ideal of R is w -stable;
- (vii) R is w -stable and each t -maximal ideal of R is t -invertible.

Proof. (i) \Rightarrow (ii). If M is a t -maximal ideal of R , then R_M is an integrally closed stable domain by Theorem 2.2. Hence R_M is a valuation domain [36, Theorem 4.6].

(ii) \Rightarrow (iii). For each t -maximal ideal M of R , R_M is a valuation stable domain (Theorem 2.2). Hence R_M is a strongly discrete valuation domain [10, Proposition 5.3.8]. The t -finite character follows again from Theorem 2.2.

(iii) \Leftrightarrow (iv) by Lemma 2.5.

(iv) \Rightarrow (v) R_M is stable, for each $M \in t\text{-Max}(R)$, because it is a strongly discrete valuation domain [10, Proposition 5.3.8]. By the t -finite character, R is w -stable (Theorem 2.2).

(v) \Rightarrow (vii) because each t -maximal ideal of a generalized Krull domain is t -invertible [6, Corollary 3.6].

(vii) \Rightarrow (i) By Theorem 2.2, R_M is a local stable domain, for each $M \in t\text{-Max}(R)$. Since M is t -invertible, MR_M is a principal ideal. Hence R_M is a valuation domain [39, Lemma 4.5] and R is integrally closed.

(iv) \Leftrightarrow (vi) follows from Theorem 2.4. □

By Theorem 2.4, each divisorial ideal of a generalized Krull domain is w -stable. Hence a w -divisorial generalized Krull domain is w -stable. Several characterizations of w -divisorial generalized Krull domains were given in [9, Theorem 3.5]. The following theorem says something more in terms of w -stability; similar results for Prüfer domains were obtained by Olberding [36, 38].

Theorem 2.7. *Let R be an integral domain. The following conditions are equivalent:*

- (i) R is an integrally closed w -divisorial w -stable domain;
- (ii) R is a w -stable w -divisorial PvMD;
- (iii) R is a w -divisorial generalized Krull domain;
- (iv) R is a weakly Matlis w -stable PvMD;
- (v) R is a weakly Matlis strongly discrete PvMD;
- (vi) R is a weakly Matlis generalized Krull domain.

Proof. (i) \Leftrightarrow (ii) by Theorem 2.6.

(ii) \Rightarrow (iv) because a w -divisorial domain is weakly Matlis [9, Theorem 1.5].

(iii) \Leftrightarrow (v) by [9, Theorem 3.5].

(v) \Leftrightarrow (vi) by Lemma 2.5.

(iii) + (vi) \Rightarrow (i) and (iv) \Leftrightarrow (v) by Theorem 2.6, because a weakly Matlis domain has t -finite character. □

From Theorems 2.6 and 2.7, we immediately get:

Corollary 2.8. *Let R be an integrally closed w -stable domain. Then R is w -divisorial if and only if each nonzero t -prime ideal is contained in a unique t -maximal ideal.*

A domain is stable and divisorial if and only if it is totally divisorial [38, Theorem 3.12]. The following is the t -analogue of this result in the integrally closed case.

Corollary 2.9. *Let R be an integral domain. The following conditions are equivalent:*

- (i) R is a w -stable w -divisorial PvMD;
- (ii) R is integrally closed and each t -linked overring of R is w -divisorial.

Proof. By [9, Theorem 3.5], if R is integrally closed, each t -linked overring of R is w -divisorial if and only if R is a weakly Matlis strongly discrete PvMD. We conclude by applying Theorem 2.7. \square

We recall that each overring of a domain R is t -linked if and only if $d = w$ on R [4, Theorem 2.6] and that each overring of a stable domain is stable [39, Theorem 5.1]. We now prove that w -stability is preserved by t -linked extension.

Theorem 2.10. *Let R be an integral domain and T a t -linked overring of R . If R is w -stable then T is w' -stable, where w' denotes the w -operation on T .*

Proof. We shall use Theorem 2.2. Since $R \subseteq T$ is t -linked, for each t' -maximal ideal M of T , there is a t -maximal ideal N of R such that $R_N \subseteq T_M$ [4, Proposition 2.1]. Hence T_M is an overring of a stable domain and is therefore stable [39, Theorem 5.1].

We next show that T has t -finite character. Let N be a t -maximal ideal of R and let $\{M_\alpha\}$ be a family of t -maximal ideals of T such that $\bigcap_\alpha M_\alpha \neq (0)$ and $M_\alpha \cap R \subseteq N$ for each α . Set $S := \bigcap_\alpha T_{M_\alpha}$. Then S is a stable domain since it is an overring of the stable domain R_N [39, Theorem 5.1]. The prime ideals $P_\alpha = M_\alpha T_{M_\alpha} \cap S$ of S are pairwise incomparable, since $S_{P_\alpha} = T_{M_\alpha}$ for each α . We have $(0) \neq \bigcap_\alpha M_\alpha \subseteq \bigcap_\alpha P_\alpha$, and, since S is treed [39, Theorem 4.11 (ii)] and has finite character [39, Theorem 3.3], then $\{P_\alpha\}$ must be finite. Hence $\{M_\alpha\}$ is also a finite set. Since R has t -finite character, it follows that T has t -finite character. \square

We do not know whether the integral closure of a w -stable domain is w' -stable. In fact the integral closure of a domain R is not always t -linked over R [5, Section 4] and we cannot apply Theorem 2.10. However, the w -integral closure $R^{[w]} := \bigcup\{(J_w : J_w); J \text{ a finitely generated ideal of } R\}$ is integrally closed and t -linked over R [4, Proposition 2.2 (a)]. Thus we immediately get:

Corollary 2.11. *The w -integral closure of a w -stable domain is a w' -stable PvMD.*

We end by remarking that, in the integrally closed case, w -divisoriality and w -stability correspond to divisoriality and stability of the t -Nagata ring. We shall make use of Proposition 1.3.

Theorem 2.12. *Let R be an integral domain. Then:*

- (1) *R has t -finite character if and only if $R\langle X \rangle$ has finite character.*
- (2) *R is a Weakly Matlis PvMD if and only if $R\langle X \rangle$ is an h -local Prüfer domain.*
- (3) *R is a strongly discrete PvMD if and only if $R\langle X \rangle$ is a strongly discrete Prüfer domain.*
- (4) *R is a generalized Krull domain if and only if $R\langle X \rangle$ is a generalized Dedekind domain.*
- (5) *R is an integrally closed w -divisorial domain if and only if $R\langle X \rangle$ is an integrally closed divisorial domain.*
- (6) *R is an integrally closed w -stable domain if and only if $R\langle X \rangle$ is an integrally closed stable domain.*

Proof. Denote by $c(f)$ the content of a polynomial $f(X) \in R[X]$.

(1) We have $\text{Max}(R\langle X \rangle) = \{MR\langle X \rangle; M \in t\text{-Max}(R)\}$. Since $f(X) \in MR[X]$ if and only if $c(f)_v \subseteq M$, if R has t -finite character, then $R\langle X \rangle$ has t -finite character. The converse is clear.

(2) Follows from (1) and Proposition 1.3.

(3) For $M \in t\text{-Max}(R)$, we have that $R\langle X \rangle_{MR\langle X \rangle} = R[X]_{MR[X]} = R_M(X)$ is a strongly discrete valuation domain if and only if R_M has the same property.

(4) Follows from (3) and Proposition 1.3 by recalling the definitions.

(5) When R is integrally closed, R is divisorial if and only if R is an h -local Prüfer domain such that each maximal ideal is invertible [23, Theorem 5.1] and R is w -divisorial if and only if R is a weakly Matlis PvMD such that each t -maximal ideal is t -invertible [9, Theorem 3.3]. Hence we can

conclude by applying part (2) and recalling that, for each $M \in t\text{-Max}(R)$, $MR\langle X \rangle$ is invertible if and only if M is t -invertible [30, Theorem 2.4].

(6) Follows from [36, Theorem 4.6], Theorem 2.6 and statements (1) and (3). \square

REFERENCES

- [1] D. D. Anderson and M. Zafrullah, *Independent locally-finite intersections of localizations*, Houston J. Math. **25** (1999), 433-452.
- [2] H. Bass, *On the ubiquity of Gorenstein rings*, Math. Z. **82** (1963), 8-28.
- [3] S. Bazzoni and L. Salce, *Warfield Domains*, J. Algebra **185** (1996), 836-868.
- [4] D.E. Dobbs, E.G. Houston, T.G. Lucas, and M. Zafrullah, *t -linked overrings and Prüfer v -multiplication domains*, Comm. Algebra, **17** (1989), 2835-2852.
- [5] D.E. Dobbs, E.G. Houston, T.G. Lucas, M. Roitman and M. Zafrullah, *On t -linked overrings*, Comm. Algebra **20** (1992), 1463-1488.
- [6] S. El Baghdadi, *On a class of Prüfer v -multiplication domains*, Comm. Algebra **30** (2002), 3723-3742.
- [7] S. El Baghdadi, *Factorization of divisorial ideals in a generalized Krull domain*, Rings, Modules, Algebras and Abelian Groups (Venice, 2002), 149-160, Lecture notes in Pure and appl. Math. **236**, Marcel Dekker, New York, 2004.
- [8] S. El Baghdadi and M. Fontana, *Semistar linkedness and flatness, Prüfer semistar multiplication domains*, Comm. Algebra **32** (2004), 1101-1126.
- [9] S. El Baghdadi and S. Gabelli, *w -divisorial domains*, J. Algebra **285** (2005), 335-355.
- [10] M. Fontana, J. Huckaba, and I. Papick, *Prüfer domains*, Dekker, New York, 1997.
- [11] M. Fontana, P. Jara and E. Santos, *Prüfer \star -multiplication domains and semistar operations*, J. Algebra Appl. **2** (2003) 21-50.
- [12] M. Fontana and K. A. Loper, *Nagata rings, Kronecker function rings and related semistar operations*, Comm. Algebra **31** (2003), 4775-4805.
- [13] S. Gabelli, *A class of Prüfer domains with nice divisorial ideals*, Commutative Ring Theory (Fés, 1995), 313-318, Lecture Notes in Pure and Appl. Math. **185**, M. Dekker, New York, 1997.
- [14] S. Gabelli, *Prüfer ($\#\#$)-domains and localizing systems of ideals*, Advances in Commutative Ring Theory (Fés, 1997), 391-409, Lecture Notes in Pure and Appl. Math. **205**, M. Dekker, New York, 1999.
- [15] S. Gabelli, E. Houston and T. Lucas *The $t\#$ -property for integral domains*, J. Pure Appl. Algebra, **194** (2004), 281-298.
- [16] S. Gabelli and G. Picozza, *Star stable domains*, manuscript.
- [17] R. Gilmer, *Integral domains which are almost Dedekind*, Proc. Am. Math. Soc. **15** (1964), 813-818.
- [18] R. Gilmer, *Multiplicative ideal theory*, Dekker, New York, 1972.
- [19] R. Gilmer, *Overrings of Prüfer domains*, J. Algebra **4** (1966), 331-340.
- [20] R. Gilmer and W. Heinzer, *Overrings of Prüfer domains. II*, J. Algebra **7** (1967), 281-302.
- [21] R. Gilmer and W. Heinzer, *Primary ideals with finitely generated radical in a commutative ring*, Manuscripta Math. **78** (1993), 201-221.
- [22] M. Griffin, *Some results on Prüfer v -multiplication rings*, Canad. J. Math. **19** (1967), 717-722.
- [23] W. J. Heinzer, *Integral domains in which each non-zero ideal is divisorial*, Matematika **15** (1968), 164-170.
- [24] W. J. Heinzer and I. J. Papick, *The radical trace property*, J. Algebra **112** (1988), 110-121.
- [25] E. Houston and M. Zafrullah, *Integral domains in which each t -ideal is divisorial*, Michigan Math. J. **35** (1988), 291-300.
- [26] E. Houston and M. Zafrullah, *On t -invertibility II*, Comm. Algebra **17** (1989), 1955-1969.
- [27] E. Houston, S. Kabbaj, T. Lucas and A. Mimouni, *When is the dual of an ideal a ring?*, J. Algebra **225** (2000), 429-450.
- [28] J. A. Huckaba and I. J. Papick, *Connecting trace properties*, Non-Noetherian Commutative Ring Theory, 313-324, Math. Appl. **520**, Kluwer Acad. Publ., Dordrecht, 2000.
- [29] B.G. Kang, *\star -Operation on integral domains*, Ph.D. Thesis, The University of Iowa, 1987.
- [30] B.G. Kang, *Prüfer v -multiplication domains and the ring $R[X]_{N_v}$* , J. Algebra **123** (1989), 151-170.
- [31] D.J. Kwak and Y.S. Park, *On t -flat overrings*, Ch. J. Math. **23** (1995), 17-24.
- [32] J. Lipman, *Stable ideals and Arf rings* J. Pure Appl. Algebra **4** (1974), 319-336.
- [33] T. Lucas, *The radical trace property and primary ideals*, J. Algebra **184** (1996), 1093-1112.
- [34] E. Matlis, *Reflexive domains*, J. Algebra **8** (1968), 1-33.

- [35] A. Mimouni, *Star-operations and trace properties*, International Journal of Commutative Rings, **1** (2002), 77-92.
- [36] B. Olberding, *Globalizing local properties of Prüfer domains*, J. Algebra **205** (1998), 480-504.
- [37] B. Olberding, *On the classification of stable domains*, J. Algebra **243** (2001), 177-197.
- [38] B. Olberding, *Stability, duality, 2-generated ideals and a canonical decomposition of modules*, Rend. Sem. Mat. Univ. Padova **106** (2001), 261-290.
- [39] B. Olberding, *On the structure of stable domains*, Comm. Algebra **30** (2002), 877-895.
- [40] N. Popescu, *On a class of Prüfer domains*, Rev. Roumanie Math. Pures Appl. **29** (1984), 777-786.
- [41] J.D. Sally and W.V. Vasconcelos, *Stable rings and a problem of Bass*, Bull. Am. Math. Soc. **79** (1973), 575-576.
- [42] J.D. Sally and W.V. Vasconcelos, *Stable rings*, J. Pure Appl. Alg. **40** (1974), 319-336.
- [43] Wang Fanggui and R.L. McCasland, *On w -modules over Strong Mori domains*, Comm. Algebra **25** (1997), 1285-1306.
- [44] Wang Fanggui and R.L. McCasland, *On Strong Mori domains*, J. Pure Appl. Algebra **135** (1999), 155-165.

DEPARTMENT OF MATHEMATICS, FACULTÉ DES SCIENCES ET TECHNIQUES, P.O. BOX 523, BENI MELLAL, MOROCCO

E-mail address: `baghdadi@fstbm.ac.ma`

DIPARTIMENTO DI MATEMATICA, UNIVERSITÀ DEGLI STUDI "ROMA TRE", LARGO S. L. MURIALDO, 1, 00146 ROMA, ITALY

E-mail address: `gabelli@mat.uniroma3.it`