

# Elementary Number Theory (TN410)

## Exercises: Sheet #1

March 4, 2016

1. Compute  $\gcd(5520, 3135)$ ,  $\gcd(8736, 3135)$ ;
2. Compute  $v_2(70!)$ ,  $v_5(125!)$  and  $v_7(130!)$ ;
3. Let  $a, b, c, n \in \mathbb{N}$ . Show that
  - (a) If  $a \mid n$ ,  $b \mid n$  and  $\gcd(a, b) = 1$ , then  $ab \mid n$
  - (b) If  $a \mid bc$  and  $\gcd(a, b) = 1$ , then  $a \mid c$ .
4. Show that there exist infinitely many primes  $p$  of the form  $p = 4k - 1$ ;  
(*hint*: Assume that  $p_1, \dots, p_s$  are the only primes of this form and consider  $N = 4p_1 \cdots p_s - 1$ )
5. Let  $\pi(x) = \#\{p \leq x\}$ .
  - (a) Compute (by hand or with a computer)  $\pi(10)$ ,  $\pi(100)$ ,  $\pi(1000)$  and  $\pi(10000)$ ;
  - (b) Compare, in each case, the obtained value both with  $X/\log X$  and with  $\text{li}(X)$ .
6. Let, for  $k > 1$ ,  $\zeta(k) = \sum_{n=1}^{\infty} \frac{1}{n^k}$ . Show that

$$\sum_{n \leq X} \frac{1}{n^k} = \zeta(k) + O\left(\frac{1}{X^{k-1}}\right) \quad \text{and that} \quad \sum_{n \leq X} \frac{\mu(n)}{n^k} = \frac{1}{\zeta(k)} + O\left(\frac{1}{X^{k-1}}\right);$$

7. We say that  $n \in \mathbb{N}$  is  $k$ -free if, for each prime  $p$ ,  $p^k \nmid n$ . Let  $\mu_k$  be the characteristic function of  $k$ -free integers. that is:

$$\mu_k(n) = \begin{cases} 1 & \text{if } n \text{ is } k\text{-free;} \\ 0 & \text{otherwise.} \end{cases}$$

- (a) Show that  $\mu_k$  is multiplicative;
- (b) Prove the identity:

$$\mu_k(n) = \sum_{\substack{d \in \mathbb{N} \\ d^k \mid n}} \mu(d);$$

- (c) Show that

$$\sum_{n \leq X} \mu_k(n) = \frac{1}{\zeta(k)} X + O(X^{1/k}).$$

8. Show that the probability that two positive integers are coprime, is  $6/\pi^2$ ;
9. Let  $N$  be an hypothetical odd perfect number. Show that the unique factorization of  $N$  has the form:

$$N = p_1^k \cdot p_2^{2j_2} \cdots p_r^{2j_r}$$

where  $k \geq 1$ ,  $j_1, \dots, j_r \geq 1$  and  $p_1 \equiv k \equiv 1 \pmod{4}$ ;

(*hint*: note that it must be  $\sigma(N) = 2N \equiv 2 \pmod{4}$  and deduce from it some properties of  $\sigma(p^\alpha)$  for  $p^\alpha \parallel N$ )

10. Show that an odd perfect number  $N$  cannot be of the form  $6m - 1$ .